

Isotopic production cross-sections and recoil velocities of spallation-fission fragments in the reaction $^{238}\text{U}(1\text{A GeV})+\text{d}$

J. Pereira,^{1,*} J. Benlliure,¹ E. Casarejos,¹ P. Armbruster,² M. Bernas,³ A. Boudard,⁴ S. Czajkowski,⁵ T. Enqvist,² R. Legrain,⁴ S. Leray,⁴ B. Mustapha,³ M. Pravikoff,⁵ F. Rejmund,³ K.-H. Schmidt,² C. Stéphan,³ J. Taïeb,³ L. Tassan-Got,³ C. Volant,⁴ and W. Wlazlo⁴

¹*Universidad de Santiago de Compostela, Santiago de Compostela E-15782, Spain*

²*Gesellschaft für Schwerionenforschung, Darmstadt D-64291, Germany*

³*Institute de Physique Nucléaire, Orsay cedex F-91406, France*

⁴*DAPNIA/SPH, CEA/Saclay, Gif sur Yvette cedex F-91191, France*

⁵*CENBG, Gradignan cedex F-33175, France*

(Dated: April 30, 2006)

Fission fragments of 1A GeV ^{238}U nuclei interacting with a deuterium target have been investigated with the Fragment Separator (FRS) at GSI (Darmstadt) by measuring their isotopic production cross-sections and recoil velocities. The results, along with those obtained recently for spallation-evaporation fragments, provide a comprehensive analysis of the spallation nuclear productions in the reaction $^{238}\text{U}(1\text{A GeV})+\text{d}$. Details about the experimental performance, data reduction and results will be presented.

PACS numbers: 21.10.Ft; 21.10.Gv; 25.40.Sc; 25.45.-z; 25.75.-q; 25.85.-w; 29.25.Rm

I. INTRODUCTION

Spallation reactions constitute nowadays one of the most extensively used processes to produce Radioactive Nuclear Beams (RNB's) upon the entire chart of the nuclides [1]. Besides their numerous technical and scientific applications (for a review see for instance Refs. [2–6]), this type of reactions has recently turned out to be an optimum tool for investigating the de-excitation of hot nuclei over a wide range of temperatures and fissilities [6–10].

The present paper focuses on the experimental study of the spallation-fission fragments produced in the reaction ^{238}U at 1A GeV with deuterium, and constitutes the complementary part of the analysis of spallation-evaporation fragments presented recently by E.Casarejos et al. [11]. Both experiments are enclosed within an extensive experimental campaign performed at GSI to determine the general inventory of nuclear fragments produced in different spallation reactions [6, 12–23]. The main goal of the project was the establishment of an extensive benchmark data basis for the design of Accelerator-Driven Systems (ADS) [24, 25] and spallation-neutron sources. A restrictive requirement of these applications is the highly-precise measurement of the isotopic production cross-sections and recoil velocities of spallation-fragments.

The challenging performance of this kind of experiments motivated the use of the inverse-kinematics technique, in which the heavy-ion accelerated at relativistic energies collides with a light-nucleus target in such a

way that the spallation-reaction products are emitted forward allowing thus their analysis with an in-flight magnetic separator. At present, the only laboratory offering the possibility of employing such a technique for studying uranium-induced reactions at relativistic energies the GSI [26] in Darmstadt, Germany. Details about this facility, as well as the experimental technique used to separate and identify the reaction products are presented in the first part of the paper. The second part concerns the data analysis performed to evaluate the recoil velocities and isotopic production cross-sections of the measured fission fragments. The paper ends with a compilation and general discussion of the results. An extensive analysis of these results will be presented in a forthcoming paper.

II. EXPERIMENT

A. Experimental setup

In the present experiment, a ^{238}U primary beam was accelerated in the heavy-ion synchrotron SIS [27, 28] up to an energy of 1A GeV. The intensity, of the order of 10^7 ions/s, was continuously measured during the experiment with the secondary-electron monitor SEETRAM [29, 30]. After its extraction, the uranium beam impinged on a cryogenic deuterium target [31] of 201 mg/cm² thickness, isolated from the vacuum line by two titanium foils of 18.16 mg/cm² thickness each. This target was located at 1.69 m from the entrance of the Fragment Separator (FRS) [32], so that the resulting reaction products were transmitted and measured by the separator before their β -decay. A niobium stripper foil of 60 mg/cm² thick was placed downstream the production target in order to maximize the fraction of fully stripped

*Electronic address: pereira@nscsl.msu.edu; Present address: NSCL/MSU, East Lansing MI-48824, US




FIG. 1: Schematic view of the FRS with its standard detection equipment.

fragments.

The FRS is an achromatic zero-degree in-flight separator, which consists of four independent stages with their respective image planes (see Fig. 1). The four stages are grouped in two symmetric dispersive areas, so that the entire device remains achromatic. Its high resolving power of 1500, determined for an emittance of 20π mm-mrad and a beam spot of 2.7 mm [32], is achieved at the expenses of a limited acceptance in magnetic rigidity ($\pm 1.5\%$) and angle (15 mrad around the beam axis). According to the ion-optics of the FRS, the separation and identification of the fission fragments was performed by combining the signals measured with the standard detection setup shown in Fig. 1:

A pair of plastic scintillators (*SC2* and *SC4* [33]), placed at the second (*F2*) and fourth (*F4*) image planes were used to determine the horizontal (x direction) deflection of the transmitted nuclei and the time-of-flight from *F2* to *F4*. Two ionization chambers (*MUSIC's* [34]), placed at the end of the FRS, provided the energy loss of the nuclei. Finally, two multiwire chambers (*MW41* and *MW42* [35]) were used to track the transmitted nuclei. All the fast-output signals were filtered by setting the CFD threshold-levels above the electronic noise and the low-amplitude signals generated by light-particles. The trigger for the data acquisition was given by the signals from the last plastic scintillator (*SC4*). In order to consider a valid event, a software-defined trigger condition required valid position, time-of-flight, angle and energy loss signals.

B. Separation and identification of reaction products by the FRS

1. Separation of transmitted nuclei

The magnetic separation of the nuclides transmitted through the FRS was based upon the relationship between the mass-over-charge ratio A/Z of the nuclei, their

magnetic rigidity $B\rho$, and the longitudinal component of the relativistic reduced momentum $\beta\gamma$:

$$B\rho = \left(\frac{A}{Z}\right) \cdot \frac{m_0}{e} \cdot \beta\gamma \quad (1)$$

being m_0 the nuclear mass unit and e the electron charge.

For a given nucleus, its magnetic rigidity was determined from the horizontal deflection, measured with the plastic scintillator *SC2* at the dispersive image plane, according to:

$$B\rho = (B\rho)_0 \cdot \left(1 - \frac{x_2}{D_2}\right) \quad (2)$$

where x_2 is the position of the transmitted nucleus, measured in the dispersive plane *F2*; D_2 is the dispersion at this image plane, corresponding to the first half of the separator; and $(B\rho)_0$ corresponds to the magnetic rigidity of a charged particle following a central trajectory along the FRS. This latter quantity was determined by multiplying the magnetic fields of each dipole, measured with Hall probes, with the corresponding curvature radii.

The longitudinal component of the reduced momentum $\beta\gamma$ of the transmitted nuclei (or equivalently their longitudinal velocity v) was calculated by dividing the path length l covered by the ions from *F2* to *F4* by the time-of-flight *ToF* between the two plastic scintillators *SC2* and *SC4*. In order to avoid dependences of the velocity on the angle, an angular correction from the tracking detectors *MW41* and *MW42* had to be included in the path-length l .

Together with the mass-over-charge ratio A/Z , we also determined the atomic number of each nuclide, in order to get a full identification. This quantity was deduced from the energy-loss signal ΔE measured with the ionization chambers (*MUSIC's*), corrected for the dependencies on the velocity of the transmitted nuclei and on the horizontal positions of the incoming nuclei at the entrance of the detector with respect to the anodes.

The final resolution attained in the separation of the reaction products is illustrated in the left panel of Fig. 2, where the energy-loss signal is plotted against the mass-over-charge ratio for some selected nuclei. The figure was obtained by summing up all the fission settings measured during the experiment.

As far as the selection of transmitted nuclei is concerned, the transversal dimensions of the inner tubes of the FRS and the maximum magnetic dispersion set a limit on the range of magnetic rigidities accepted by the separator. Assuming a maximum dispersion of -6.8 cm/% at the dispersive image plane, a $B\rho$ -acceptance of $\pm 1.5\%$ was deduced from Eq. 2; those nuclei with magnetic rigidities 1.5% greater or lesser than the central value $(B\rho)_0$ were deflected out of the FRS beam line. This limited range of accepted $B\rho$'s, together with the velocity spread, defined a region of A/Z values that could be measured in one single tuning (setting) of the magnetic dipoles.




FIG. 2: Two-dimensional cluster plots showing the energy-loss signal against the mass-over-charge ratio. Left: Separated nuclei, represented by the different blobs. Right: Isospin identification (the vertical line corresponds to nuclei with $N=Z$)

Since the nuclei produced in the collision of ^{238}U at 1A GeV with the deuterium target spread out over a wide region of atomic and mass numbers, the FRS had to be adjusted according to the magnetic rigidities of the medium-mass fission fragments to be analyzed. From the calculated values of $B\rho$, a central value $(B\rho)_0$ of the FRS was selected by properly adjusting the magnetic fields of each dipole magnet according to the curvature radii. In order to cover the region of interest, the measurements were repeated for 33 settings centered on different values of $(B\rho)_0$, separated from each other in steps of 1%.

2. Isotopic identification

The identification of the atomic number Z for each nucleus was deduced using the energy-loss signals (ΔE) of the projectile-nucleus as a reference.

The identification of mass numbers was obtained from the matrix $(A/Z) - Z$ (Fig. 2, right): depending on the values of the mass and atomic numbers of the nucleus, the different nuclei are located at different blobs in this matrix. The possible combinations obtained between the quantities A/Z and Z —where Z and A are integer numbers—defined a characteristic pattern in which the nuclei could be grouped within curved lines. The only line with no curvature corresponds to light nuclei with the same number of protons and neutrons ($Z = N$). The next group-line on the right corresponds to nuclei with $N = Z + 1$, which differs a bit from a perfect vertically. If one then goes n steps to the right, the line found corresponds to the group of isotopes with $N = Z + n$. Thus, from the vertical group-line $A = 2Z$ it was possible to identify the mass number of all the nuclides, since their atomic number was well known. The same rule applied

to the next lines to the right ($N = Z + 1, N = Z + 2, \dots$) in such a way that all the nuclei could be identified in mass A and atomic Z numbers.

Figure 3 shows the identification matrix obtained by summing up all the different settings measured during the experiment. From this figure, about 1000 different nuclides could be identified.

III. DATA REDUCTION

A. Determination of recoil velocities

The recoil velocities of the fission fragments were evaluated in order to investigate the kinematics associated with the reaction mechanism. The spallation reaction itself is characterized by the collision of a heavy uranium projectile with a deuterium target nucleus, followed by the de-excitation of the projectile-like fragments by fission or by evaporation of light particles. The kinematics of the final fragments correspond to the convolution of the velocity distribution produced by the fast collision, and the subsequent de-excitation processes.

In the present experiment, the kinematics of the fission and evaporation fragment productions were separated by measuring the velocity distribution along the beam axis (hereafter referred to as v), filtered by the limited angular acceptance of the FRS. As discussed in Refs. [18, 36], this limitation truncates the velocity distributions of fission and evaporation fragments in a different manner, in such a way that they could be identified from the resulting measured v distribution. By making use of Eq. 2, it was possible to determine the longitudinal velocities of the nuclei transmitted through the FRS from their positions at the dispersive image plane (F2) and the identified




FIG. 3: Identification matrix in form of a scatter plot, showing the atomic number versus the mass-over-charge of all the nuclei measured in 33 different settings of the reaction $^{238}\text{U}(1\text{A GeV})+\text{d}$. As an example, the fission fragment ^{130}Sn is indicated.

values of Z and A :

$$v = c \cdot \left\{ \left[\frac{A/Z \cdot m_0 c / e}{(B\rho)_0 \cdot \left(1 - \frac{x_2}{D_2}\right)} \right]^2 + 1 \right\}^{-1/2} \quad (3)$$

where c refers to the speed of light. In order to determine the exact recoil velocities of the reaction products, the values obtained from Eq. 3 had to be corrected by the energy losses of the transmitted nuclei in the different layers of matter situated between the deuterium target and the plastic scintillator at F2 (see fig. 1). This correction was done by calculating the energy loss of each nucleus along the flight path from the middle of the tar-

get —where the reaction was assumed to take place on average— to the second image plane. Finally, the corrected velocities were transformed into the frame of the projectile.

The accuracy of the velocity determined from Eq. 3 was mainly limited by the position resolution of the plastic scintillator $SC2$, $\Delta x_2 \simeq 2\text{ mm}$ and the relative resolution of $(B\rho)_0$, of the order of 10^{-4} . These values led to a final relative precision of the velocity of about 3×10^{-4} which proves to be a real achievement when compared with other experimental techniques.

In order to reconstruct the v distribution of a given nucleus, the contributions measured in different magnetic settings of the FRS were summed up and normalized to




FIG. 4: Scatter plots of ${}_{40}\text{Zr}$, ${}_{46}\text{Pd}$, ${}_{60}\text{Nd}$ and ${}_{65}\text{Tb}$ elements showing the velocity distributions measured in the frame of the projectile as a function of the mass number.

the corresponding beam intensities N_{proj} . As an example of this procedure, Fig. 4 depicts the reconstructed velocity spectra for different nuclei as a function of the mass number in form of a scatter plot. The upper and lower wings correspond to the transmitted forward- and backward-emitted fission fragments, respectively, while the central cloud represents the contribution of evaporation. The centroid of this latter distribution is slightly slowed down with respect to the velocity of the projectile, while the two external wings approach each other as the fragment charge increases. The separation between these two wings is related to the fission velocity v_{fiss} and its reduction can be partially attributed to a decrease of the kinetic energy of the two fission partners due to momentum conservation in mass-asymmetric systems and, to a lesser degree, to the weakness of the Coulomb repulsion at large charge-asymmetries. Apart from this, the distribution of fragments produced in the two reaction mechanisms is consistent with theoretical expectations: fission fragments are mostly produced with greater neutron excess, while evaporation generally leads to nuclei

along the so-called evaporation corridor, located on the neutron-deficient side of the β -stability valley. Furthermore, as the charge of the element increases, the evaporation contribution is enhanced until it becomes the dominant production mechanism.

The velocities of the fission fragments v_{fiss} in the frame of the projectile were obtained for each isotope by fitting the measured velocity spectra to specific functions which reproduced the contribution from each reaction mechanism. Figure 5 depicts an example of the fit obtained for ${}^{83}\text{Sr}$. The central peak due to the evaporation component, was described by a Gaussian function $f_{evap}(v)$, while the forward and backward fission peaks were best reproduced by fitting functions defined by the convolution of a Gaussian and an exponential function. Each of these functions $f_{fiss}^f(v)$ and $f_{fiss}^b(v)$ had four free parameters, three of which corresponded to the Gaussian components, while the fourth parameter represented the skewness of the inner tails. The spectra were fitted to a global function $F(v)$ with eleven parameters (three from the Gaussian evaporation component and eight from the




FIG. 5: Velocity-spectrum fit for ^{83}Sr showing the contributions from evaporation (central Gaussian curve) and fission (external curves) to the global fit.

forward and backward fission fit-functions):

$$F(v) = f_{fiss}^b(v) + f_{evap}(v) + f_{fiss}^f(v) \quad (4)$$

The spectra of nuclei produced only by fission or evaporation were well described by the functions of the specific component, only.

As can be seen in Fig. 6, the radius of the spherical fission distribution, corresponding to the velocity of the fission fragments v_{fiss} , could be obtained by halving the distance between the external sides of the forward and backward fission peaks, given by v_f and v_b :

$$v_{fiss} = \frac{|v_f - v_b|}{2} = \frac{|v_f| + |v_b|}{2} \quad (5)$$

However, the determination of v_f and v_b from the velocity spectra was quite complicated due to the smooth side tails of the fission peaks.

Alternatively, the values of v_f and v_b could be expressed as a function of the most probable velocities v_f^{max} and v_b^{max} that corresponded to the maximum of the forward and backward fitting functions $f_f(v)$ and $f_b(v)$, according to:

$$v_f = v_f^{max} + \frac{v_{fiss} - v_{fiss} \cos \alpha_f}{2} \quad (6)$$

and

$$v_b = v_b^{max} + \frac{v_{fiss} + v_{fiss} \cos \alpha_b}{2} \quad (7)$$

where α_f and α_b are the forward and backward angles obtained by a Lorentz transformation of the angular acceptance α_{eff} of the FRS into the frame of the fissioning system [36, 37]. These two equations can be re-written as:

$$v_f = v_f^{max} + v_{fiss} \cdot T_f(v_{fiss}) \quad (8)$$




FIG. 6: Schematic representation of the velocity distribution of fission fragments (solid circumference) affected by the cuts produced by the limited angular acceptance of the FRS (dashed lines). (Note the shift of the velocity distribution with respect to the velocity of the projectile, marked by the origin of the reference frame). The projection of the transmitted nuclei onto the axis parallel to the beam direction (longitudinal component) is shown on the lower part of the figure. The measured most-probable forward and backward fission velocities v_f^{max} and v_b^{max} are compared with the real values v_f and v_b .

and

$$v_b = v_b^{max} + v_{fiss} \cdot T_b(v_{fiss}) \quad (9)$$

where $T_f(v_{fiss})$ and $T_b(v_{fiss})$ are the v_{fiss} -dependent transmissions of the forward- and backward-emitted fission fragments, calculated according to Ref. [36].

The idea behind Eqs. 8 and 9 was that the velocities v_f^{max} and v_b^{max} , obtained from the fitting functions, must be corrected using the fission velocity v_{fiss} and the angular transmission, in order to provide the values v_f and v_b . Because v_{fiss} depends on v_f and v_b (see Eq. 5), and T_f and T_b on v_{fiss} , we used the recursive algorithm described in Ref. [36] to deduce both the forward- and backward-transmissions T_f , T_b , and the fission velocity v_{fiss} . The iterative calculation of the algorithm stops when the measured velocities v_f^{max} and v_b^{max} converge to constant values, which are assumed to correspond to v_f and v_b .

There were two sources of uncertainty in the above-described method: first was the determination of v_f^{max} and v_b^{max} from fits and second was the uncertainty of the angular acceptance value α_{eff} used to calculate T_f and T_b . The uncertainties of the former varies considerably from one nucleus to another and depends mainly on the smoothness of the velocity profile; it ranged typically from 1% to 5%, reaching in some particular cases maximum values up to 20%. Concerning the uncertainty due to α_{eff} , it mainly affected the velocities of fragments with rather low transmission. A careful analysis of the ion-optics and the geometry of the FRS yielded a total

uncertainty for α_{eff} of about 4% [36], resulting in an uncertainty for the velocities of less than 5%.

Finally, an additional source of error arose from the production of secondary reactions in the deuterium target induced by a heavy primary fission fragment. Such a process would reduce the atomic and mass numbers of the fission fragments, while slightly blurring its velocity spectra. As a consequence, the velocity distribution of the isotope can be partially contaminated by the velocities of the heavier fission fragments that underwent a secondary reaction, thus leading to a reduction of the measured values of v_f^{max} and v_b^{max} , as shown in Ref. [18]. This contamination was only significant for the most neutron-deficient fragments.

B. Determination of production cross-sections

In the thin-target approximation, the total production yield of a given isotope is defined by:

$$Y_{tot}(Z, A) = \frac{N(Z, A)}{N_{proj}} \quad (10)$$

where $N(Z, A)$ is the number of nuclei, with a given Z and A , measured during the experiment, and N_{proj} is the number of beam particles. The value of $N(Z, A)$ was obtained by integrating the measured velocity spectra for each nucleus. These spectra were partly contaminated by the fragment production on the titanium windows surrounding the cryogenic deuterium target. In order to subtract the contamination from the reactions produced in the titanium windows, all the measurements were repeated with a titanium dummy target of thickness equivalent to the target windows. The total yields $Y_{tot}(Z, A)$, integrated over the isotopic chains, are compared in Fig. 7 with the contribution from the titanium dummy target (or equivalently the titanium windows) Y_{dummy} . The production cross-sections for the reaction $^{238}\text{U}+d$ were determined from:

$$\sigma_{prod}(Z, A) = \frac{1}{\chi} \cdot [Y_{tot}(Z, A) - Y_{dummy}(Z, A)] \cdot F \quad (11)$$

being χ the density of deuterium nuclei integrated over the target thickness. The factor F includes the different corrections that had to be applied to the measured data due to the limitations of the experimental setup. This factor was calculated according to:

$$F = f_{eff} \cdot f_{sr} \cdot f_{ch} \cdot f_{ev} \cdot f_{\tau} \quad (12)$$

where f_{eff} accounts for the efficiencies of the experimental setup, f_{sr} corresponds to the corrections that were necessary due to secondary reactions, f_{ch} considers the charge states of the measured nuclei, f_{ev} corrects the contamination from evaporation reactions, and f_{τ} takes into account the dead time of the data acquisition. The calculation of these correction factors has been extensively




FIG. 7: Element distribution of fission-fragment production yields in the cryogenic deuterium target Y_{tot} (dots) and the titanium dummy target Y_{dummy} (triangles).

described in previous analysis of fission- and evaporation-fragment productions (see for instance Refs. [6, 12–23]). Consequently, in the present discussion we rather restrict ourselves to mention their main features.

The correction due to limitations of the efficiency f_{eff} included the intrinsic efficiency of the detection setup and the angular transmission of the FRS. The former was evaluated for each element by comparing the total number of valid events with the number of events detected with the *MUSIC*'s detectors (with an efficiency of 100%). The resulting values were above 98% in most of the cases. The correction due to the limited angular transmission of the FRS was evaluated according to the method described in Ref. [36]. This method was based on the calculation of the angular acceptance α_{eff} and the analysis of the fission kinematics. The resulting total angular transmission varied largely from values near 100%, for heavy fragments with rather narrow angular emittance, to values around 15%, for the lightest nuclei. The uncertainty associated with this correction method arose from two different sources, namely the calculation of the angular acceptance α_{eff} and the analysis of the kinematics. The final transmission correction error varied significantly from about 2% to 30%, for some particular cases.

Apart from the primary reactions in the target, the beam particles and the reaction products could undergo further reactions in the different layers of matter placed along the FRS, thereby modifying the primary fragment productions. The list of possible “parasitic” targets included the accelerator window, the aluminum SEETRAM foils, the titanium windows surrounding the cryogenic target, a niobium stripper foil behind the target, and the plastic scintillator *SC2*. The probability for a given nucleus to undergo a secondary reaction was estimated by multiplying the production cross sections, ob-




FIG. 8: Measured velocity spectra of three cesium isotopes measured in the reaction $^{238}\text{U}(1\text{A GeV})+\text{d}$.

tained from Glauber-type model calculations [38, 39], by the number of impinging beam particles and the number of target nuclei of the corresponding layer of matter. Most of the fragments produced in the aluminum SEETRAM foils and the accelerator window were not transmitted through the FRS due to their large distance with respect to the entrance of the separator that reduced significantly the value of α_{eff} . Moreover, since these contaminating targets were also present in the settings with the titanium dummy target, their contribution to the production cross sections were already subtracted according to eq. 11. The correction due to secondary reactions induced in the niobium foil and the plastic scintillator *SC2* reached rather low values ranging from 10%, for the lightest elements to almost 20% for the heaviest. Furthermore, the primary fragments produced in the reaction $^{238}\text{U}(1\text{A GeV})+\text{d}$ could undergo a secondary reaction before leaving the rather thick deuterium target. These secondary products could also induce a ternary reaction resulting in a multi-step reaction chain. The multiple reaction contaminants did not spread uniformly over the total fragment production but concentrated in some specific regions close to the evaporation corridor. In the case of fission fragments, this contamination mainly affected the neutron-deficient isotopes, gradually increasing as the neutron number decreases. In the present analysis, we used the correction method proposed by P. Napolitani et al. [40]. According to these authors, the measured productions (referred to as apparent cross-sections $\tilde{\sigma}$) were formulated through a system of master equations that included the losses of fragments due to their interaction in the target and the gains due to the contamination from intermediate multiple reactions. A detailed discussion of this model and the underlying approximations used to solve the set of master equations can be found in Ref. [40]. For each given element, the correction factor was found to be constant within 5% to 7% for most of the isotopes, and to drastically drop down when approaching the neutron-deficient side, wherein the corrections reached values close to 100%.

Besides secondary nuclear reactions, the residual nu-

clei could also pick-up or strip-off some electrons due to the atomic interactions with the different layers of matter. The corresponding fraction of bare ions was maximized by placing a niobium stripper foil of 60 mg/cm^2 thickness behind the target. However, the charge-state configuration of the transmitted ions could change from the first to the second stage of the FRS due to the interactions with the plastic scintillator *SC2*. As explained in Ref. [41], the strong correlation between the charge state of the transmitted ions and their transversal positions at the final image plane makes it possible to discriminate any charge-state change along the FRS. In order to avoid ambiguities in the charge identification, we restricted the present analysis to ions with the same charge-state configuration in both stages. Besides bare ions, this includes hydrogen-like and helium-like ions; though their extremely low fraction (less than 1%) for the range of charges covered by this analysis did not affect the final result. According to this criterion, the losses of measured nuclei which were not fully stripped in the two stages of the FRS had to be corrected. For each nucleus, the correction factor f_{ch} was calculated with the GLOBAL code [42, 43] as the inverse of the probability to be fully stripped along the FRS, i.e. the product of the fractions of bare ions behind the niobium stripper foil (first stage) and behind the scintillator *SC2* (second stage), respectively. The resulting correction factor kept always below 10% within the range of elements analyzed in the present work.

Regarding the correction due to the dead time of the data acquisition f_{τ} , the total number of physical triggers generated by the logic of the experiment and the number of events processed by the data acquisition were counted for each setting with scalers. The ratio between both numbers provided the value of the dead time, which was around 15% for most of the settings. Furthermore, the intensity of the primary beam was checked continuously in order to keep the dead time below 30%.

Finally, since the aim of the present work was the determination of the production cross-sections of fission fragments, we had to disentangle the evaporation-




FIG. 9: Velocity distributions as a function of the atomic number of fragments measured with the deuterium target enclosed in the titanium container (left) and with the titanium dummy target (right).

reaction contributions to the measured cross-sections. By comparing the productions of all the fragments produced in the cryogenic deuterium target and the titanium dummy target we saw that the evaporation fragments measured in the former system were mainly produced in the titanium windows. However, after subtracting these contaminants, a small evaporation component was still observed in the reaction $^{238}\text{U}(1\text{A GeV})+d$ for heavy neutron-deficient isotopes. In a former analysis of spallation-evaporation fragments produced in this reaction [11], it was found that these nuclei populated the evaporation corridor situated on the neutron-deficient side of the chart of the nuclides, from uranium isotopes down to elements with atomic numbers about $Z=58$. Below this value, the evaporation component dropped quickly –though still contributing to the production of the most neutron-deficient isotopes– whereas the fission production increased very fast, specially in the neutron-rich side. Therefore, we expected both reaction mechanisms to coexist in the region of light evaporation fragments, with a transition from evaporation to fission as the neutron number increases. Figure 8 illustrates this trend by showing the velocity spectra of three different cesium isotopes. The lightest and heaviest isotopes were mainly produced by evaporation and fission, respectively, while an intermediate situation was found for ^{125}Cs , wherein a weak evaporation component was strongly mixed with the dominant fission mechanism (see also Ref. [11]).

The evaporation contamination to the measured fission fragments was estimated using a phenomenological model based on the two-step formalism [44]. A Glauber-type Monte-Carlo subroutine [15] was chosen to reproduce the fast interaction between the projectile and target. The evaporation stage was simulated with the statistical de-excitation ABLA code [10, 39] and the PROFI fission mass-distribution subroutine [45]. Details of these codes are well beyond the scope of the present paper; it is sufficient to mention here that the reliability of the calcula-

tions in estimating evaporation productions in the region of neutron-deficient heavy nuclei was verified by comparing their results with the evaporation cross-sections measured for some nuclei. Our estimations agreed with the experimental results within the precision requirements of the present analysis. Furthermore, the separation method discussed in reference [11] was used in the present analysis for some nuclei and compared with the technique described above. Both methods agreed within their respective uncertainties, thus demonstrating their capabilities to disentangle the two reaction mechanisms.

IV. RESULTS

A. Velocities of fission fragments

Figure 9 shows the velocity distributions of the fission fragments (in the projectile frame) measured in the reaction induced in the deuterium target surrounded by the titanium windows (left) and in the titanium dummy target (right). The distributions were obtained for each element by integrating the contributions of the whole isotopic chain. In the figure, the evaporation fragments lay in a region of velocities close to that of the primary beam, while the forward- and backward-emitted fission fragments correspond to the upper and lower wings. The increase of the fission velocities as the atomic number of the fragments decreases is a natural consequence of the momentum conservation between the light and heavy fission fragments in the frame of the fissioning system.

The velocities v_{fiss} of the fission fragments with atomic numbers ranging from $Z = 23$ to $Z = 66$ were determined according to the method described in section III. Figures 10-12 show the isotopic chains of these velocities v_{fiss} for all the measured fragments. The numerical values, with their associated uncertainties are included in Tabs. I-VIII.




FIG. 10: Isotopic dependence of velocities of fission fragments produced in $^{238}\text{U}(1\text{A GeV})+\text{d}$ (dots) compared to the values calculated with the scission-model [46] (see text for details). The calculations were obtained for different fissioning systems: uranium Z=92 (solid line), radium Z=88 (dashed line), mercury Z=80 (dotted line) and rhenium Z=75 (dash-dotted line). Empty dots correspond to nuclei with a secondary-reaction contamination greater than 50%. (Total errors are shown when larger than symbols).




FIG. 11: Isotopic dependence of velocities of fission fragments produced in $^{238}\text{U}(1\text{A GeV})+\text{d}$ (dots) compared to the values calculated with the scission-model [46] (see text for details). The calculations were obtained for different fissioning systems: uranium $Z=92$ (solid line), radium $Z=88$ (dashed line), mercury $Z=80$ (dotted line) and rhenium $Z=75$ (dash-dotted line). Empty dots correspond to nuclei with a secondary-reaction contamination greater than 50%. (Total errors are shown when larger than symbols).




FIG. 12: Isotopic dependence of velocities of fission fragments produced in $^{238}\text{U}(1\text{A GeV})+\text{d}$ (dots) compared to the values calculated with the scission-model [46] (see text for details). The calculations were obtained for different fissioning systems: uranium $Z=92$ (solid line), radium $Z=88$ (dashed line), mercury $Z=80$ (dotted line) and rhenium $Z=75$ (dash-dotted line). Empty dots correspond to nuclei with a secondary-reaction contamination greater than 50%. (Total errors are shown when larger than symbols).

As a general trend, one observes a general reduction of the average fission velocities as the charge of the fission fragments increases. This trend is explained by momentum conservation of the two fission partners: after scission, the total kinetic energy of the system (i.e. the sum of the kinetic energies of each fission partner) comes almost entirely from the Coulomb repulsion between the two nascent fragments. Due to momentum conservation, the lightest nucleus will acquire more velocity than the heavier one, thus explaining the results shown in the figures.

Regarding the trends observed for a given element the velocities reach a maximum value for the most neutron-rich isotopes, then evolve through a plateau for less neutron-rich nuclei and finally decrease in the region of light isotopes. This final reduction is more pronounced in the most neutron deficient side with an abrupt decline of v_{fiss} to values 20% below the maximum. The characteristic pattern shown in these figures was already observed in the analysis of fission fragments produced in the reaction $^{238}\text{U}(1\text{A GeV})+\text{p}$ [18]. According to this work, the abrupt decrease of the velocities in the neutron-deficient side can be attributed, to some extent, to the contamination of secondary reactions from primary fission fragments. As discussed above, in order to estimate such an effect, we have used the model proposed by P. Napolitani et al. [40]. Those nuclei with a secondary-reaction component greater than 50% are represented in Figs. 10-12 by empty dots. As expected, they correspond to the most neutron-deficient isotopes.

In order to understand the observed isotopic dependence of the velocities, a comparison between the values measured in the present experiment and those obtained from the scission-model developed by Wilkins et al. [46] is shown in Figs. 10-12. The different lines drawn on top of the experimental values correspond to velocities acquired by fission isotopes coming from different fissioning systems. Surprisingly, the distribution of these systems

extends over a rather wide range of elements: the most neutron-rich isotopes stem from the heaviest fissioning elements including uranium, while the neutron-deficient ones are populated by lighter fissioning systems like mercury. This trend is in agreement with the Z_{fiss}^2/A_{fiss} -dependence of the fissility: As a consequence of the low fissilities of light parent nuclei (low Z_{fiss}), only the most neutron-deficient isotopes (low A_{fiss}) have a chance for undergoing fission due to their smaller fission barriers; the residues produced by these parent nuclei populate thus the neutron-deficient side. On the contrary, the high fissilities of heavy fissioning systems are rather insensitive to a variation of the neutron number, thus extending their productions to more neutron-rich nuclei. A detailed analysis of these results will be discussed in a forthcoming paper.

B. Isotopic production cross-sections of fission fragments

The production cross-sections of 939 fission fragments measured in the present work are represented as a scatter plot on the top of the chart of the nuclides in Fig. 13 (see also Tabs. IX-XVII). They cover a region on the neutron-rich side of the valley of stability, with elements ranging from vanadium to thulium. The most populated region corresponds to elements between niobium and cadmium with decreasing intensities for lower and higher atomic numbers. For most of these elements, the maximum productions span an extended flat ridge shifted by 2-3 mass units to the right side of the valley of stability. Above the doubly closed shell $Z=50$ $N=82$, an enhanced fission-fragment production is observed on the neutron-rich side.

Figures 14-16 provide a more detailed survey of the fission-fragment production, showing the isotopic production cross sections for all measured fission fragments. The measured cross sections range from about 30 mb for




FIG. 13: (Colored on-line) Fission fragments in the reaction $^{238}\text{U} + \text{d}$ at 1A GeV on top of the chart of the nuclides. The colored scale indicates different production cross-sections.

intermediate masses of the different elements, down to values as low as $2 \mu\text{b}$ on the neutron-rich side.

As a general trend, the isotopic chains of the fission fragments follow a wide Gaussian distribution with mean neutron numbers slightly shifted toward the neutron-rich side of the valley of stability. The maximum production covers an extended region of elements around ^{46}Pd , which corresponds to symmetric fission. For some elements above $Z=50$, these Gaussian shapes display a pronounced shoulder on the neutron-rich side related to the contribution of the asymmetric fission channels standard I and standard II [47] by low-energy fission of nuclei close to ^{238}U . The polarization of the heavy asymmetric fission fragments toward the neutron-rich side, makes the lighter asymmetric partners to move toward the valley of stability, leading to a less pronounced shoulder near the maximum of the isotopic chains, as shown for elements between ^{33}As and ^{42}Mo . This double-humped structure is not observed for elements above ^{59}Pr and below ^{30}Zn .

The total fission cross-section, obtained by summing up all the individual isotopic production cross-sections was found to be $2.00 \pm 0.42 \text{ b}$. In order to account for the isotopes which were not measured, we extrapolated the isotopic chains shown in Figs. 14-16. The contribution from the missed residues to the total fission cross-section was found to be negligible compared with the total uncertainty. At the present moment, fission data of uranium on deuterium are scarcely represented

in the current available databases. After a systematic literature search on the EXFOR [48] database, the only measurements found corresponded to deuteron-induced fission of uranium at energies below 200 MeV [49]. A comparison of these data with our results was possible by extrapolating the former according to the systematics of Prokofiev [50] for proton-induced fission reactions. In doing so, we had to consider the two main differences between proton and deuteron reactions at these energies, namely the average double energy introduced by deuterons, due to their additional nucleon, and the higher total reaction cross-section with respect to protons. Thus, the total fission cross-section measured by Stevenson et al. [49] at 190 MeV for deuteron-induced reactions on uranium ($2.49 \pm 0.05 \text{ b}$) was compared to the value given by Prokofiev at $2 \times 190 \text{ MeV} = 380 \text{ MeV}$ for protons (1.44 b). The differences between these data were entirely attributed to the higher total reaction cross-section for deuterium, leading to a scaling factor of $S = 2.49/1.44 = 1.73$. In order to extrapolate the value of Stevenson measured at 190 MeV to 1 GeV, we multiplied the cross-section calculated by Prokofiev at 2 GeV (1.18 b) by the factor S , which yields 2.04 b. This value compares very well with our result.

The total reaction cross-sections could be determined, as well by adding the total evaporation cross-section of $0.91 \pm 0.11 \text{ b}$ obtained by E. Casarejos [11] for the reaction $^{238}\text{U}(1\text{A GeV}) + \text{d}$ to the total fission cross-section

measured in the present work. The resulting value of 2.91 ± 0.43 b, is still compatible with the 2.51 b calculated with the Glauber-type model of P.J. Karol [38], which assumes a Gaussian density profile with $r(\text{rms})=2.1\text{fm}$ for the deuteron nuclei [51]

In contrast to other existing experimental techniques, the total fission cross section obtained in the present work was not a directly measured observable, but deduced by summing up the contributions from every fission fragment. The isotopic production cross-sections of the fission fragments shown in Figs. 14-16 were obtained by applying different systematic corrections to the measured yields. Some of these corrections, like the angular transmission or the secondary reactions in the target, vary considerably as a function of the atomic and mass numbers of the final nuclei, reaching very large values in some cases. Any uncertainty in the determination of these corrections affected the values of the isotopic cross-sections, and consequently the value of the total cross-section. The high level of agreement between the total fission cross-section obtained in the present work and that deduced by extrapolation of a directly measured value [49] validates the methods used to determine these corrections. The 15% difference between the measured total reaction cross-section and that calculated with Karol's model is within the uncertainty of our method.

V. CONCLUSION

In the present work, the spallation-fission fragments produced in the reaction $^{238}\text{U}(1\text{A GeV})$ on deuterium were experimentally investigated. The results obtained from the analysis include the evaluation of the velocities and isotopic production cross-sections of 939 fission nuclei, which, together with the 602 evaporation fragments measured recently [11], constitutes one of the most extensive inventories of spallation-reaction productions ever measured before. The elements included in both experiments spread out over an extend region in the chart of the nuclides, ranging from vanadium to thulium for the fission case, and from cerium to neptunium for evaporation.

The precisions required for the present measurements were only attainable with the heavy-ion experimental facility at GSI, Darmstadt (Germany). The uranium beam, accelerated in the Synchrotron accelerator SIS up to 1A GeV, was focused onto a cryogenic deuterium target placed at the entrance of the Fragment Separator (FRS), so that the forward-emitted fission fragments were transmitted through the separator. The optimum performance of the FRS, with a momentum resolving power of 1500, allowed the challenging separation of the reaction products, as well as their unambiguous identification according to the atomic and mass numbers.

The velocity distributions of the fission fragments were measured with relative resolutions of the order of 10^{-4} . Fits to these velocity distributions allowed for the sep-

aration of the kinematics from evaporation and fission components. The results from the fits were introduced in an algorithm to extract the velocities of the fission fragment. The precisions attained with this technique ranged typically from 1% to 5%, reaching maximum values of 20% in cases of very poor statistics or high contamination from evaporation reactions. Within a given element, the isotopic dependence of the fission velocities revealed a contribution from many different fissioning systems to the total productions. Moreover, the relatively low velocities of some neutron-deficient isotopes were found to be a sign of the contamination from multiple reactions induced by primary fission fragments.

The isotopic production cross-sections of the fission fragments were measured above the lower limit of $2\mu\text{b}$, with statistical uncertainties ranging from 1% to 5% for the majority of the measured fragments. The systematic error induced by the different corrections applied to the measured isotopic productions covers a wide range which extends from 10-20%, for most of the nuclei, to maximum values of 50% for very few cases. The main source of systematic error were the corrections that accounted for the multiple reactions in the production target and for the contamination from evaporation reactions. The maximum value of the cross-sections –around 30 mb– were found for intermediate-mass fragments around palladium. The isotopic chains of most of the fission fragments are well described by a Gaussian function with mean neutron numbers slightly shifted towards the neutron-rich side of the valley of stability. For some elements above $Z=50$, a second maximum shows up in the neutron-rich isotopes, revealing a sizeable population of the asymmetric fission channels by low-energy fission. When summing up the measured isotopic cross-sections, a total value of 2.00 ± 0.42 b was found, which is in good agreement with results obtained from Glauber-type models and extrapolations from experimental values. The addition of this cross-section to the value of 0.91 ± 0.11 b for the evaporation component [11], yields a total value of 2.91 ± 0.43 b, which is in good agreement with Glauber-type calculations [38].

The richness of the present results demands a detailed discussion of the many different physical aspects underlying the investigated reaction.

Acknowledgments

This work was partially supported by the Spanish Ministry of Education and Science and Xunta de Galicia under contracts FPA2002-04181-C04-01 and PGIDT01PXI20603PM, respectively, and the European Community under contracts “Access to Research Infrastructure Action of the Improving Human Potential” PRIC-1999-00001, “HINDAS” FIKW-CT-2000-00031 and “Research Infrastructure Action - Structuring the European Research Area” EURISOL DS Project Contract no. 515768 RIDS.




FIG. 14: Measured isotopic production cross-sections of fission fragments produced in the reaction $^{238}\text{U}(1\text{A GeV})+\text{d}$. (Statistical errors are shown when larger than symbols).




FIG. 15: Measured isotopic production cross-sections of fission fragments produced in the reaction $^{238}\text{U}(1\text{A GeV})+\text{d}$. (Statistical errors are shown when larger than symbols).




FIG. 16: Measured isotopic production cross-sections of fission fragments produced in the reaction $^{238}\text{U}(1\text{A GeV})+\text{d}$. (Statistical errors are shown when larger than symbols).

The authors wish to thank K.-H. Behr, A. Brüne and K. Burkard for their technical support during the experiment, as well as the group of P. Chesny, who built the liquid-hydrogen/deuterium target.

The EC is not liable for any use that may be made of the information contained herein.

APPENDIX A: VELOCITY AND ISOTOPIC PRODUCTION CROSS-SECTIONS OF FISSION FRAGMENTS

-
- [1] RNB VI Nucl. Phys. A748 (2005)
 - [2] The European Spallation Source Study, The ESS Technical Study, Vol. III, Report ESS-96-53-M, 1996
 - [3] C.D. Bowman, E.D. Arthur, P.W. Lisowski, G.P. Lawrence, R.J. Jensen, J.L. Anderson, B. Blind, M. Cappiello, J.W. Davidson, T.R. England, L.N. Engel, R.C. Gaight, H.G. Hughes, J.R. Ireland, R.A. Krakowski, R.J. LaBauve, B.C. Letellier, R.T. Perry, G.J. Russel, K.P. Staudhammer, G. Versamis, and W.B. Wilson, Nucl. Instr. and Meth. A320 (1992) 336
 - [4] W.F. Henning, Nucl. Instr. and Meth. B126 (1997) 1
 - [5] R. Michel, I. Leya, and L. Borges, Nucl. Instr. and Meth. B113 (1996) 343
 - [6] J. Pereira, Ph.D. Thesis, Universidade de Santiago de Compostela (2004)
 - [7] J. Benlliure, P. Armbruster, M. Bernas, A. Boudard, T. Enqvist, R. Legrain, S. Leray, F. Rejmund, K.-H. Schmidt, C. Stéphan, L. Tassan-Got, and C. Volant, Nucl. Phys. A700 (2002) 469
 - [8] P. Napolitani, K.-H. Schmidt, A.S. Botvina, F. Rejmund, L. Tassan-Got, and C. Villagrasa, Phys. Rev. C70 (2004) 054607
 - [9] K.-H. Schmidt, M.V. Ricciardi, A. Botvina, and T. Enqvist, Nucl. Phys. A710 (2002) 157
 - [10] A.R. Junghans, M. de Jong, H.-G. Clerc, A.V. Ignatyuk, G.A. Kudyaev and K.-H. Schmidt, Nucl. Phys. A629 (1998) 635
 - [11] E. Casarejos, J. Benlliure, J. Pereira, P. Armbruster, M. Bernas, A. Boudard, S. Czajkowski, T. Enqvist, R. Legrain, S. Leray, B. Mustapha, M. Pravikoff, F. Rejmund, K.-H. Schmidt, C. Stéphan, J. Taïeb, L. Tassan-Got, C. Volant, and W. Wlazlo, submitted to Phys. Rev. C
 - [12] W. Wlazlo, T. Enqvist, P. Armbruster, J. Benlliure, M. Bernas, A. Boudard, C. Czajkowski, R. Legrain, S. Leray, B. Mustapha, M. Pravikoff, F. Rejmund, K.-H. Schmidt, C. Stéphan, J. Taïeb, L. Tassan-Got, and C. Volant, Phys. Rev. Lett. 84 (2000) 5736
 - [13] J. Benlliure, P. Armbruster, M. Bernas, A. Boudard, J.P. Dufour, T. Enqvist, R. Legrain, S. Leray,

nucleus	v_{fiss} (cm/ns)	ε_{tot} (cm/ns)	nucleus	v_{fiss} (cm/ns)	ε_{tot} (cm/ns)
^{49}V	1.590	0.070	^{65}Co	1.620	0.051
^{50}V	1.690	0.020	^{66}Co	1.620	0.051
^{51}V	1.720	0.022	^{67}Co	1.620	0.051
^{52}V	1.700	0.010	^{68}Co	1.620	0.081
^{53}V	1.750	0.030	^{69}Co	1.620	0.081
^{54}V	1.760	0.010	^{57}Ni	0.870	0.150
^{55}V	1.710	0.040	^{58}Ni	1.030	0.120
^{56}V	1.710	0.100	^{59}Ni	1.250	0.071
^{57}V	1.710	0.100	^{60}Ni	1.400	0.020
^{49}Cr	0.840	0.150	^{61}Ni	1.470	0.032
^{50}Cr	1.140	0.120	^{62}Ni	1.530	0.010
^{51}Cr	1.450	0.090	^{63}Ni	1.570	0.014
^{52}Cr	1.630	0.050	^{64}Ni	1.590	0.010
^{53}Cr	1.700	0.010	^{65}Ni	1.610	0.010
^{54}Cr	1.690	0.010	^{66}Ni	1.630	0.010
^{55}Cr	1.720	0.010	^{67}Ni	1.600	0.014
^{56}Cr	1.690	0.010	^{68}Ni	1.600	0.071
^{57}Cr	1.690	0.050	^{69}Ni	1.620	0.060
^{58}Cr	1.690	0.060	^{70}Ni	1.640	0.022
^{59}Cr	1.690	0.080	^{71}Ni	1.620	0.070
^{51}Mn	0.880	0.160	^{72}Ni	1.620	0.100
^{52}Mn	1.110	0.130	^{59}Cu	1.010	0.160
^{53}Mn	1.520	0.120	^{60}Cu	1.150	0.120
^{54}Mn	1.660	0.050	^{61}Cu	1.340	0.100
^{55}Mn	1.670	0.010	^{62}Cu	1.450	0.061
^{56}Mn	1.710	0.022	^{63}Cu	1.530	0.022
^{57}Mn	1.690	0.014	^{64}Cu	1.550	0.014
^{58}Mn	1.650	0.010	^{65}Cu	1.570	0.014
^{59}Mn	1.690	0.010	^{66}Cu	1.550	0.014
^{60}Mn	1.650	0.014	^{67}Cu	1.560	0.010
^{61}Mn	1.650	0.022	^{68}Cu	1.580	0.014
^{62}Mn	1.650	0.051	^{69}Cu	1.600	0.010
^{63}Mn	1.640	0.051	^{70}Cu	1.550	0.030
^{53}Fe	0.860	0.150	^{71}Cu	1.550	0.040
^{54}Fe	1.160	0.100	^{72}Cu	1.610	0.060
^{55}Fe	1.450	0.070	^{73}Cu	1.610	0.070
^{56}Fe	1.570	0.022	^{74}Cu	1.610	0.071
^{57}Fe	1.650	0.070	^{62}Zn	1.010	0.120
^{58}Fe	1.620	0.010	^{63}Zn	1.210	0.091
^{59}Fe	1.610	0.010	^{64}Zn	1.370	0.022
^{60}Fe	1.610	0.020	^{65}Zn	1.430	0.014
^{61}Fe	1.620	0.050	^{66}Zn	1.520	0.010
^{62}Fe	1.600	0.051	^{67}Zn	1.520	0.014
^{63}Fe	1.600	0.051	^{68}Zn	1.540	0.010
^{64}Fe	1.600	0.061	^{69}Zn	1.550	0.010
^{65}Fe	1.600	0.041	^{70}Zn	1.530	0.010
^{66}Fe	1.600	0.041	^{71}Zn	1.560	0.010
^{55}Co	0.820	0.150	^{72}Zn	1.570	0.020
^{56}Co	1.030	0.120	^{73}Zn	1.570	0.040
^{57}Co	1.340	0.081	^{74}Zn	1.570	0.040
^{58}Co	1.540	0.010	^{75}Zn	1.570	0.014
^{59}Co	1.600	0.030	^{76}Zn	1.580	0.040
^{60}Co	1.630	0.030	^{77}Zn	1.580	0.050
^{61}Co	1.630	0.030	^{78}Zn	1.580	0.050
^{62}Co	1.620	0.050	^{64}Ga	0.960	0.110
^{63}Co	1.620	0.051	^{65}Ga	1.210	0.081
^{64}Co	1.620	0.051	^{66}Ga	1.320	0.071

TABLE I: Velocities of spallation-fission fragments (in cm/ns) measured in the frame of the fissioning system for the reaction $^{238}U(1A\text{ GeV})+d$, with their total error.

nucleus	v_{fiss} (cm/ns)	ε_{tot} (cm/ns)	nucleus	v_{fiss} (cm/ns)	ε_{tot} (cm/ns)
^{67}Ga	1.440	0.032	^{79}Se	1.470	0.010
^{68}Ga	1.470	0.010	^{80}Se	1.470	0.010
^{69}Ga	1.510	0.010	^{81}Se	1.460	0.014
^{70}Ga	1.530	0.014	^{82}Se	1.460	0.014
^{71}Ga	1.520	0.014	^{83}Se	1.450	0.032
^{72}Ga	1.520	0.014	^{84}Se	1.450	0.041
^{73}Ga	1.540	0.014	^{85}Se	1.450	0.014
^{74}Ga	1.540	0.014	^{86}Se	1.480	0.014
^{75}Ga	1.540	0.030	^{87}Se	1.490	0.010
^{76}Ga	1.540	0.030	^{88}Se	1.490	0.030
^{77}Ga	1.540	0.032	^{89}Se	1.480	0.041
^{78}Ga	1.540	0.032	^{73}Br	1.110	0.091
^{79}Ga	1.540	0.051	^{74}Br	1.200	0.071
^{80}Ga	1.540	0.061	^{75}Br	1.290	0.060
^{67}Ge	1.090	0.120	^{76}Br	1.310	0.030
^{68}Ge	1.300	0.080	^{77}Br	1.360	0.014
^{69}Ge	1.400	0.061	^{78}Br	1.400	0.014
^{70}Ge	1.440	0.040	^{79}Br	1.410	0.014
^{71}Ge	1.490	0.010	^{80}Br	1.420	0.014
^{72}Ge	1.520	0.010	^{81}Br	1.410	0.022
^{73}Ge	1.490	0.014	^{82}Br	1.410	0.020
^{74}Ge	1.520	0.014	^{83}Br	1.450	0.010
^{75}Ge	1.520	0.014	^{84}Br	1.450	0.010
^{76}Ge	1.510	0.010	^{85}Br	1.450	0.050
^{77}Ge	1.540	0.010	^{86}Br	1.450	0.030
^{78}Ge	1.520	0.050	^{87}Br	1.470	0.010
^{79}Ge	1.520	0.050	^{88}Br	1.470	0.010
^{80}Ge	1.500	0.014	^{89}Br	1.470	0.010
^{81}Ge	1.510	0.014	^{90}Br	1.450	0.010
^{82}Ge	1.520	0.014	^{91}Br	1.450	0.030
^{83}Ge	1.520	0.041	^{92}Br	1.450	0.040
^{69}As	1.100	0.090	^{76}Kr	1.150	0.071
^{70}As	1.270	0.050	^{77}Kr	1.230	0.061
^{71}As	1.340	0.030	^{78}Kr	1.280	0.032
^{72}As	1.410	0.022	^{79}Kr	1.330	0.010
^{73}As	1.420	0.014	^{80}Kr	1.340	0.010
^{74}As	1.430	0.014	^{81}Kr	1.370	0.010
^{75}As	1.480	0.030	^{82}Kr	1.370	0.014
^{76}As	1.460	0.022	^{83}Kr	1.400	0.014
^{77}As	1.480	0.010	^{84}Kr	1.400	0.014
^{78}As	1.480	0.010	^{85}Kr	1.410	0.014
^{79}As	1.490	0.010	^{86}Kr	1.400	0.014
^{80}As	1.490	0.040	^{87}Kr	1.400	0.014
^{81}As	1.490	0.030	^{88}Kr	1.420	0.040
^{82}As	1.490	0.010	^{89}Kr	1.420	0.010
^{83}As	1.470	0.010	^{90}Kr	1.430	0.014
^{84}As	1.470	0.010	^{91}Kr	1.420	0.010
^{85}As	1.490	0.014	^{92}Kr	1.400	0.010
^{86}As	1.500	0.050	^{93}Kr	1.430	0.014
^{71}Se	1.060	0.091	^{94}Kr	1.410	0.014
^{72}Se	1.220	0.071	^{95}Kr	1.410	0.041
^{73}Se	1.280	0.010	^{78}Rb	1.070	0.100
^{74}Se	1.330	0.010	^{79}Rb	1.180	0.070
^{75}Se	1.390	0.014	^{80}Rb	1.280	0.040
^{76}Se	1.430	0.010	^{81}Rb	1.310	0.022
^{77}Se	1.440	0.010	^{82}Rb	1.330	0.014
^{78}Se	1.450	0.010	^{83}Rb	1.340	0.010

TABLE II: Velocities of spallation-fission fragments (in cm/ns) measured in the frame of the fissioning system for the reaction $^{238}U(1A\text{ GeV})+d$, with their total error.

nucleus	v_{fiss} (cm/ns)	ε_{tot} (cm/ns)	nucleus	v_{fiss} (cm/ns)	ε_{tot} (cm/ns)
⁸⁴ Rb	1.370	0.010	⁸⁵ Zr	1.090	0.071
⁸⁵ Rb	1.380	0.022	⁸⁶ Zr	1.110	0.051
⁸⁶ Rb	1.370	0.010	⁸⁷ Zr	1.170	0.030
⁸⁷ Rb	1.380	0.010	⁸⁸ Zr	1.190	0.030
⁸⁸ Rb	1.370	0.014	⁸⁹ Zr	1.210	0.010
⁸⁹ Rb	1.400	0.032	⁹⁰ Zr	1.230	0.014
⁹⁰ Rb	1.420	0.030	⁹¹ Zr	1.260	0.014
⁹¹ Rb	1.400	0.041	⁹² Zr	1.260	0.014
⁹² Rb	1.400	0.022	⁹³ Zr	1.260	0.010
⁹³ Rb	1.420	0.014	⁹⁴ Zr	1.280	0.014
⁹⁴ Rb	1.420	0.010	⁹⁵ Zr	1.290	0.015
⁹⁵ Rb	1.400	0.020	⁹⁶ Zr	1.290	0.014
⁹⁶ Rb	1.390	0.020	⁹⁷ Zr	1.310	0.020
⁹⁷ Rb	1.370	0.020	⁹⁸ Zr	1.300	0.035
⁸⁰ Sr	1.100	0.071	⁹⁹ Zr	1.290	0.041
⁸¹ Sr	1.160	0.061	¹⁰⁰ Zr	1.310	0.032
⁸² Sr	1.190	0.061	¹⁰¹ Zr	1.320	0.030
⁸³ Sr	1.240	0.030	¹⁰² Zr	1.340	0.014
⁸⁴ Sr	1.280	0.010	¹⁰³ Zr	1.340	0.014
⁸⁵ Sr	1.310	0.014	¹⁰⁴ Zr	1.340	0.014
⁸⁶ Sr	1.320	0.010	¹⁰⁵ Zr	1.350	0.032
⁸⁷ Sr	1.320	0.010	⁸⁷ Nb	1.060	0.080
⁸⁸ Sr	1.350	0.010	⁸⁸ Nb	1.100	0.051
⁸⁹ Sr	1.360	0.014	⁸⁹ Nb	1.140	0.014
⁹⁰ Sr	1.360	0.014	⁹⁰ Nb	1.160	0.010
⁹¹ Sr	1.380	0.010	⁹¹ Nb	1.220	0.032
⁹² Sr	1.380	0.010	⁹² Nb	1.220	0.014
⁹³ Sr	1.380	0.040	⁹³ Nb	1.240	0.010
⁹⁴ Sr	1.380	0.030	⁹⁴ Nb	1.230	0.014
⁹⁵ Sr	1.390	0.020	⁹⁵ Nb	1.260	0.010
⁹⁶ Sr	1.390	0.010	⁹⁶ Nb	1.270	0.014
⁹⁷ Sr	1.390	0.014	⁹⁷ Nb	1.270	0.010
⁹⁸ Sr	1.370	0.015	⁹⁸ Nb	1.290	0.010
⁹⁹ Sr	1.350	0.017	⁹⁹ Nb	1.290	0.014
¹⁰⁰ Sr	1.350	0.020	¹⁰⁰ Nb	1.290	0.014
⁸² Y	1.010	0.150	¹⁰¹ Nb	1.290	0.032
⁸³ Y	1.140	0.091	¹⁰² Nb	1.270	0.032
⁸⁴ Y	1.200	0.050	¹⁰³ Nb	1.290	0.014
⁸⁵ Y	1.210	0.030	¹⁰⁴ Nb	1.290	0.014
⁸⁶ Y	1.230	0.020	¹⁰⁵ Nb	1.300	0.015
⁸⁷ Y	1.260	0.014	¹⁰⁶ Nb	1.300	0.020
⁸⁸ Y	1.270	0.014	¹⁰⁷ Nb	1.290	0.030
⁸⁹ Y	1.310	0.010	⁸⁹ Mo	1.010	0.091
⁹⁰ Y	1.310	0.010	⁹⁰ Mo	1.050	0.051
⁹¹ Y	1.340	0.010	⁹¹ Mo	1.110	0.041
⁹² Y	1.340	0.014	⁹² Mo	1.160	0.010
⁹³ Y	1.330	0.014	⁹³ Mo	1.190	0.010
⁹⁴ Y	1.360	0.041	⁹⁴ Mo	1.190	0.014
⁹⁵ Y	1.320	0.032	⁹⁵ Mo	1.220	0.014
⁹⁶ Y	1.340	0.041	⁹⁶ Mo	1.220	0.014
⁹⁷ Y	1.350	0.040	⁹⁷ Mo	1.230	0.014
⁹⁸ Y	1.340	0.022	⁹⁸ Mo	1.250	0.010
⁹⁹ Y	1.360	0.010	⁹⁹ Mo	1.260	0.010
¹⁰⁰ Y	1.340	0.015	¹⁰⁰ Mo	1.280	0.014
¹⁰¹ Y	1.340	0.015	¹⁰¹ Mo	1.260	0.010
¹⁰² Y	1.330	0.015	¹⁰² Mo	1.270	0.010
⁸⁴ Zr	0.940	0.100	¹⁰³ Mo	1.270	0.022

TABLE III: Velocities of spallation-fission fragments (in cm/ns) measured in the frame of the fissioning system for the reaction ²³⁸U(1A GeV)+d, with their total error.

nucleus	v_{fiss} (cm/ns)	ε_{tot} (cm/ns)	nucleus	v_{fiss} (cm/ns)	ε_{tot} (cm/ns)
¹⁰⁴ Mo	1.270	0.032	¹⁰² Rh	1.110	0.014
¹⁰⁵ Mo	1.270	0.020	¹⁰³ Rh	1.150	0.010
¹⁰⁶ Mo	1.270	0.015	¹⁰⁴ Rh	1.150	0.014
¹⁰⁷ Mo	1.280	0.015	¹⁰⁵ Rh	1.170	0.020
¹⁰⁸ Mo	1.280	0.015	¹⁰⁶ Rh	1.170	0.014
¹⁰⁹ Mo	1.270	0.020	¹⁰⁷ Rh	1.170	0.010
¹¹⁰ Mo	1.280	0.020	¹⁰⁸ Rh	1.180	0.010
⁹¹ Tc	0.910	0.100	¹⁰⁹ Rh	1.200	0.014
⁹² Tc	1.020	0.061	¹¹⁰ Rh	1.200	0.014
⁹³ Tc	1.090	0.032	¹¹¹ Rh	1.200	0.030
⁹⁴ Tc	1.110	0.014	¹¹² Rh	1.200	0.030
⁹⁵ Tc	1.130	0.014	¹¹³ Rh	1.190	0.030
⁹⁶ Tc	1.140	0.014	¹¹⁴ Rh	1.190	0.020
⁹⁷ Tc	1.160	0.020	¹¹⁵ Rh	1.180	0.014
⁹⁸ Tc	1.190	0.020	¹¹⁶ Rh	1.180	0.014
⁹⁹ Tc	1.190	0.030	¹¹⁷ Rh	1.190	0.032
¹⁰⁰ Tc	1.220	0.030	¹¹⁸ Rh	1.190	0.041
¹⁰¹ Tc	1.220	0.010	⁹⁹ Pd	0.970	0.050
¹⁰² Tc	1.220	0.014	¹⁰⁰ Pd	1.000	0.030
¹⁰³ Tc	1.230	0.010	¹⁰¹ Pd	1.020	0.032
¹⁰⁴ Tc	1.240	0.014	¹⁰² Pd	1.070	0.032
¹⁰⁵ Tc	1.220	0.020	¹⁰³ Pd	1.070	0.022
¹⁰⁶ Tc	1.210	0.020	¹⁰⁴ Pd	1.090	0.022
¹⁰⁷ Tc	1.220	0.020	¹⁰⁵ Pd	1.090	0.014
¹⁰⁸ Tc	1.220	0.025	¹⁰⁶ Pd	1.110	0.014
¹⁰⁹ Tc	1.220	0.025	¹⁰⁷ Pd	1.100	0.010
¹¹⁰ Tc	1.210	0.020	¹⁰⁸ Pd	1.140	0.014
¹¹¹ Tc	1.220	0.025	¹⁰⁹ Pd	1.150	0.014
⁹³ Ru	0.990	0.100	¹¹⁰ Pd	1.140	0.022
⁹⁴ Ru	1.000	0.061	¹¹¹ Pd	1.150	0.020
⁹⁵ Ru	1.030	0.032	¹¹² Pd	1.140	0.010
⁹⁶ Ru	1.080	0.041	¹¹³ Pd	1.140	0.020
⁹⁷ Ru	1.130	0.032	¹¹⁴ Pd	1.140	0.022
⁹⁸ Ru	1.130	0.022	¹¹⁵ Pd	1.130	0.032
⁹⁹ Ru	1.140	0.014	¹¹⁶ Pd	1.140	0.022
¹⁰⁰ Ru	1.160	0.014	¹¹⁷ Pd	1.140	0.022
¹⁰¹ Ru	1.180	0.014	¹¹⁸ Pd	1.170	0.022
¹⁰² Ru	1.170	0.010	¹¹⁹ Pd	1.160	0.014
¹⁰³ Ru	1.190	0.014	¹²⁰ Pd	1.160	0.032
¹⁰⁴ Ru	1.190	0.014	¹⁰¹ Ag	0.910	0.032
¹⁰⁵ Ru	1.190	0.022	¹⁰² Ag	0.930	0.022
¹⁰⁶ Ru	1.210	0.014	¹⁰³ Ag	0.970	0.014
¹⁰⁷ Ru	1.220	0.014	¹⁰⁴ Ag	0.970	0.032
¹⁰⁸ Ru	1.220	0.022	¹⁰⁵ Ag	1.020	0.030
¹⁰⁹ Ru	1.220	0.032	¹⁰⁶ Ag	1.040	0.010
¹¹⁰ Ru	1.210	0.030	¹⁰⁷ Ag	1.080	0.022
¹¹¹ Ru	1.210	0.014	¹⁰⁸ Ag	1.090	0.014
¹¹² Ru	1.210	0.022	¹⁰⁹ Ag	1.090	0.014
¹¹³ Ru	1.220	0.014	¹¹⁰ Ag	1.100	0.014
¹¹⁴ Ru	1.220	0.014	¹¹¹ Ag	1.130	0.010
¹¹⁵ Ru	1.220	0.014	¹¹² Ag	1.130	0.014
⁹⁶ Rh	0.920	0.081	¹¹³ Ag	1.120	0.014
⁹⁷ Rh	0.970	0.051	¹¹⁴ Ag	1.130	0.014
⁹⁸ Rh	0.980	0.051	¹¹⁵ Ag	1.150	0.010
⁹⁹ Rh	1.030	0.030	¹¹⁶ Ag	1.150	0.020
¹⁰⁰ Rh	1.080	0.010	¹¹⁷ Ag	1.130	0.032
¹⁰¹ Rh	1.100	0.014	¹¹⁸ Ag	1.130	0.032

TABLE IV: Velocities of spallation-fission fragments (in cm/ns) measured in the frame of the fissioning system for the reaction ²³⁸U(1A GeV)+d, with their total error.

nucleus	v_{fiss} (cm/ns)	ε_{tot} (cm/ns)	nucleus	v_{fiss} (cm/ns)	ε_{tot} (cm/ns)
¹¹⁹ Ag	1.130	0.022	¹¹³ Sn	0.960	0.022
¹²⁰ Ag	1.140	0.032	¹¹⁴ Sn	1.010	0.014
¹²¹ Ag	1.140	0.014	¹¹⁵ Sn	1.020	0.010
¹²² Ag	1.140	0.032	¹¹⁶ Sn	1.020	0.014
¹⁰³ Cd	0.920	0.051	¹¹⁷ Sn	1.020	0.014
¹⁰⁴ Cd	0.960	0.040	¹¹⁸ Sn	1.020	0.010
¹⁰⁵ Cd	0.970	0.032	¹¹⁹ Sn	1.030	0.014
¹⁰⁶ Cd	1.000	0.020	¹²⁰ Sn	1.040	0.014
¹⁰⁷ Cd	1.000	0.010	¹²¹ Sn	1.030	0.014
¹⁰⁸ Cd	1.020	0.032	¹²² Sn	1.040	0.014
¹⁰⁹ Cd	1.030	0.010	¹²³ Sn	1.030	0.014
¹¹⁰ Cd	1.030	0.014	¹²⁴ Sn	1.020	0.032
¹¹¹ Cd	1.030	0.014	¹²⁵ Sn	1.040	0.032
¹¹² Cd	1.060	0.014	¹²⁶ Sn	1.040	0.032
¹¹³ Cd	1.060	0.014	¹²⁷ Sn	1.050	0.012
¹¹⁴ Cd	1.080	0.010	¹²⁸ Sn	1.060	0.012
¹¹⁵ Cd	1.110	0.010	¹²⁹ Sn	1.070	0.014
¹¹⁶ Cd	1.110	0.014	¹³⁰ Sn	1.070	0.014
¹¹⁷ Cd	1.130	0.010	¹³¹ Sn	1.070	0.014
¹¹⁸ Cd	1.120	0.030	¹³² Sn	1.070	0.014
¹¹⁹ Cd	1.110	0.032	¹³³ Sn	1.060	0.014
¹²⁰ Cd	1.120	0.020	¹⁰⁹ Sb	0.660	0.081
¹²¹ Cd	1.110	0.032	¹¹⁰ Sb	0.750	0.032
¹²² Cd	1.110	0.020	¹¹¹ Sb	0.780	0.032
¹²³ Cd	1.120	0.022	¹¹² Sb	0.850	0.022
¹²⁴ Cd	1.110	0.014	¹¹³ Sb	0.880	0.014
¹²⁵ Cd	1.100	0.022	¹¹⁴ Sb	0.900	0.014
¹²⁶ Cd	1.100	0.032	¹¹⁵ Sb	0.930	0.010
¹⁰⁶ In	0.800	0.081	¹¹⁶ Sb	0.970	0.014
¹⁰⁷ In	0.880	0.051	¹¹⁷ Sb	0.970	0.014
¹⁰⁸ In	0.890	0.041	¹¹⁸ Sb	1.000	0.014
¹⁰⁹ In	0.930	0.020	¹¹⁹ Sb	1.020	0.014
¹¹⁰ In	0.980	0.014	¹²⁰ Sb	1.020	0.014
¹¹¹ In	0.980	0.014	¹²¹ Sb	1.040	0.022
¹¹² In	1.030	0.010	¹²² Sb	1.020	0.032
¹¹³ In	1.050	0.014	¹²³ Sb	1.050	0.010
¹¹⁴ In	1.050	0.014	¹²⁴ Sb	1.050	0.014
¹¹⁵ In	1.070	0.014	¹²⁵ Sb	1.040	0.014
¹¹⁶ In	1.100	0.010	¹²⁶ Sb	1.040	0.014
¹¹⁷ In	1.100	0.010	¹²⁷ Sb	1.040	0.014
¹¹⁸ In	1.090	0.014	¹²⁸ Sb	1.030	0.032
¹¹⁹ In	1.100	0.014	¹²⁹ Sb	1.050	0.014
¹²⁰ In	1.100	0.010	¹³⁰ Sb	1.060	0.010
¹²¹ In	1.100	0.020	¹³¹ Sb	1.060	0.010
¹²² In	1.110	0.032	¹³² Sb	1.060	0.014
¹²³ In	1.110	0.032	¹³³ Sb	1.070	0.014
¹²⁴ In	1.100	0.022	¹³⁴ Sb	1.060	0.014
¹²⁵ In	1.090	0.022	¹³⁵ Sb	1.050	0.032
¹²⁶ In	1.110	0.020	¹¹² Te	0.720	0.051
¹²⁷ In	1.090	0.014	¹¹³ Te	0.740	0.041
¹²⁸ In	1.090	0.022	¹¹⁴ Te	0.800	0.022
¹⁰⁷ Sn	0.700	0.061	¹¹⁵ Te	0.830	0.014
¹⁰⁸ Sn	0.780	0.032	¹¹⁶ Te	0.860	0.014
¹⁰⁹ Sn	0.820	0.014	¹¹⁷ Te	0.880	0.014
¹¹⁰ Sn	0.860	0.014	¹¹⁸ Te	0.900	0.014
¹¹¹ Sn	0.910	0.014	¹¹⁹ Te	0.920	0.024
¹¹² Sn	0.950	0.014	¹²⁰ Te	0.940	0.042

TABLE V: Velocities of spallation-fission fragments (in cm/ns) measured in the frame of the fissioning system for the reaction ²³⁸U(1A GeV)+d, with their total error.

nucleus	v_{fiss} (cm/ns)	ε_{tot} (cm/ns)	nucleus	v_{fiss} (cm/ns)	ε_{tot} (cm/ns)
¹²¹ Te	0.970	0.036	¹²⁹ Xe	0.960	0.014
¹²² Te	0.990	0.010	¹³⁰ Xe	0.990	0.020
¹²³ Te	0.990	0.014	¹³¹ Xe	0.970	0.022
¹²⁴ Te	1.000	0.010	¹³² Xe	0.970	0.014
¹²⁵ Te	1.000	0.014	¹³³ Xe	0.980	0.014
¹²⁶ Te	1.010	0.014	¹³⁴ Xe	0.970	0.022
¹²⁷ Te	1.020	0.010	¹³⁵ Xe	0.980	0.014
¹²⁸ Te	1.020	0.022	¹³⁶ Xe	0.990	0.040
¹²⁹ Te	1.010	0.022	¹³⁷ Xe	0.980	0.014
¹³⁰ Te	1.000	0.041	¹³⁸ Xe	0.980	0.022
¹³¹ Te	1.010	0.041	¹³⁹ Xe	0.950	0.020
¹³² Te	1.010	0.032	¹⁴⁰ Xe	0.940	0.032
¹³³ Te	1.020	0.015	¹⁴¹ Xe	0.970	0.022
¹³⁴ Te	1.030	0.022	¹⁴² Xe	0.950	0.022
¹³⁵ Te	1.040	0.022	¹⁴³ Xe	0.940	0.041
¹³⁶ Te	1.050	0.015	¹¹⁹ Cs	0.480	0.190
¹³⁷ Te	1.050	0.015	¹²⁰ Cs	0.650	0.081
¹³⁸ Te	1.040	0.020	¹²¹ Cs	0.700	0.041
¹³⁹ Te	1.040	0.032	¹²² Cs	0.740	0.014
¹¹⁶ I	0.630	0.100	¹²³ Cs	0.790	0.022
¹¹⁷ I	0.770	0.071	¹²⁴ Cs	0.830	0.022
¹¹⁸ I	0.830	0.032	¹²⁵ Cs	0.850	0.022
¹¹⁹ I	0.880	0.041	¹²⁶ Cs	0.890	0.022
¹²⁰ I	0.910	0.032	¹²⁷ Cs	0.910	0.022
¹²¹ I	0.910	0.014	¹²⁸ Cs	0.910	0.014
¹²² I	0.940	0.022	¹²⁹ Cs	0.900	0.014
¹²³ I	0.950	0.014	¹³⁰ Cs	0.910	0.014
¹²⁴ I	0.960	0.014	¹³¹ Cs	0.920	0.014
¹²⁵ I	0.970	0.010	¹³² Cs	0.920	0.014
¹²⁶ I	0.970	0.010	¹³³ Cs	0.920	0.014
¹²⁷ I	0.960	0.010	¹³⁴ Cs	0.920	0.014
¹²⁸ I	0.970	0.014	¹³⁵ Cs	0.920	0.014
¹²⁹ I	0.990	0.014	¹³⁶ Cs	0.900	0.032
¹³⁰ I	0.990	0.014	¹³⁷ Cs	0.900	0.041
¹³¹ I	0.990	0.014	¹³⁸ Cs	0.910	0.014
¹³² I	0.980	0.014	¹³⁹ Cs	0.900	0.032
¹³³ I	1.000	0.032	¹⁴⁰ Cs	0.890	0.022
¹³⁴ I	0.980	0.032	¹⁴¹ Cs	0.910	0.014
¹³⁵ I	1.010	0.022	¹⁴² Cs	0.910	0.014
¹³⁶ I	0.990	0.014	¹⁴³ Cs	0.910	0.014
¹³⁷ I	1.020	0.020	¹⁴⁴ Cs	0.920	0.022
¹³⁸ I	0.990	0.010	¹⁴⁵ Cs	0.910	0.022
¹³⁹ I	0.990	0.014	¹⁴⁶ Cs	0.910	0.032
¹⁴⁰ I	0.990	0.022	¹²¹ Ba	0.400	0.200
¹⁴¹ I	0.990	0.051	¹²² Ba	0.530	0.090
¹¹⁷ Xe	0.540	0.130	¹²³ Ba	0.650	0.041
¹¹⁸ Xe	0.690	0.071	¹²⁴ Ba	0.700	0.022
¹¹⁹ Xe	0.740	0.041	¹²⁵ Ba	0.730	0.022
¹²⁰ Xe	0.780	0.014	¹²⁶ Ba	0.750	0.022
¹²¹ Xe	0.830	0.051	¹²⁷ Ba	0.790	0.014
¹²² Xe	0.860	0.014	¹²⁸ Ba	0.850	0.028
¹²³ Xe	0.870	0.022	¹²⁹ Ba	0.860	0.022
¹²⁴ Xe	0.900	0.014	¹³⁰ Ba	0.890	0.014
¹²⁵ Xe	0.930	0.014	¹³¹ Ba	0.880	0.022
¹²⁶ Xe	0.930	0.014	¹³² Ba	0.920	0.014
¹²⁷ Xe	0.950	0.014	¹³³ Ba	0.920	0.014
¹²⁸ Xe	0.960	0.014	¹³⁴ Ba	0.930	0.028

TABLE VI: Velocities of spallation-fission fragments (in cm/ns) measured in the frame of the fissioning system for the reaction ²³⁸U(1A GeV)+d, with their total error.

nucleus	v_{fiss} (cm/ns)	ε_{tot} (cm/ns)	nucleus	v_{fiss} (cm/ns)	ε_{tot} (cm/ns)
¹³⁵ Ba	0.910	0.020	¹⁴² Ce	0.870	0.014
¹³⁶ Ba	0.900	0.014	¹⁴³ Ce	0.880	0.014
¹³⁷ Ba	0.920	0.014	¹⁴⁴ Ce	0.860	0.014
¹³⁸ Ba	0.910	0.014	¹⁴⁵ Ce	0.860	0.022
¹³⁹ Ba	0.910	0.014	¹⁴⁶ Ce	0.860	0.022
¹⁴⁰ Ba	0.920	0.014	¹⁴⁷ Ce	0.860	0.014
¹⁴¹ Ba	0.910	0.014	¹⁴⁸ Ce	0.860	0.014
¹⁴² Ba	0.920	0.014	¹⁴⁹ Ce	0.870	0.014
¹⁴³ Ba	0.910	0.014	¹⁵⁰ Ce	0.870	0.014
¹⁴⁴ Ba	0.910	0.014	¹⁵¹ Ce	0.850	0.022
¹⁴⁵ Ba	0.890	0.014	¹⁵² Ce	0.850	0.050
¹⁴⁶ Ba	0.890	0.014	¹²⁸ Pr	0.450	0.130
¹⁴⁷ Ba	0.890	0.022	¹²⁹ Pr	0.550	0.100
¹⁴⁸ Ba	0.890	0.022	¹³⁰ Pr	0.590	0.080
¹²⁴ La	0.440	0.200	¹³¹ Pr	0.660	0.032
¹²⁵ La	0.610	0.090	¹³² Pr	0.690	0.014
¹²⁶ La	0.680	0.051	¹³³ Pr	0.720	0.014
¹²⁷ La	0.700	0.041	¹³⁴ Pr	0.740	0.014
¹²⁸ La	0.720	0.014	¹³⁵ Pr	0.750	0.022
¹²⁹ La	0.750	0.014	¹³⁶ Pr	0.750	0.014
¹³⁰ La	0.790	0.041	¹³⁷ Pr	0.780	0.014
¹³¹ La	0.810	0.014	¹³⁸ Pr	0.770	0.022
¹³² La	0.840	0.014	¹³⁹ Pr	0.790	0.014
¹³³ La	0.850	0.014	¹⁴⁰ Pr	0.810	0.014
¹³⁴ La	0.860	0.014	¹⁴¹ Pr	0.820	0.014
¹³⁵ La	0.860	0.014	¹⁴² Pr	0.820	0.022
¹³⁶ La	0.870	0.014	¹⁴³ Pr	0.840	0.014
¹³⁷ La	0.860	0.014	¹⁴⁴ Pr	0.850	0.014
¹³⁸ La	0.860	0.014	¹⁴⁵ Pr	0.860	0.014
¹³⁹ La	0.860	0.014	¹⁴⁶ Pr	0.860	0.014
¹⁴⁰ La	0.870	0.014	¹⁴⁷ Pr	0.850	0.014
¹⁴¹ La	0.860	0.032	¹⁴⁸ Pr	0.860	0.014
¹⁴² La	0.870	0.014	¹⁴⁹ Pr	0.840	0.022
¹⁴³ La	0.870	0.014	¹⁵⁰ Pr	0.850	0.014
¹⁴⁴ La	0.860	0.032	¹⁵¹ Pr	0.840	0.014
¹⁴⁵ La	0.880	0.014	¹⁵² Pr	0.850	0.010
¹⁴⁶ La	0.870	0.014	¹⁵³ Pr	0.840	0.014
¹⁴⁷ La	0.870	0.014	¹⁵⁴ Pr	0.850	0.050
¹⁴⁸ La	0.860	0.014	¹³¹ Nd	0.430	0.150
¹⁴⁹ La	0.880	0.014	¹³² Nd	0.560	0.120
¹⁵⁰ La	0.880	0.061	¹³³ Nd	0.630	0.071
¹²⁶ Ce	0.360	0.230	¹³⁴ Nd	0.670	0.022
¹²⁷ Ce	0.540	0.100	¹³⁵ Nd	0.700	0.014
¹²⁸ Ce	0.660	0.030	¹³⁶ Nd	0.720	0.036
¹²⁹ Ce	0.710	0.014	¹³⁷ Nd	0.730	0.041
¹³⁰ Ce	0.750	0.022	¹³⁸ Nd	0.740	0.022
¹³¹ Ce	0.780	0.022	¹³⁹ Nd	0.740	0.014
¹³² Ce	0.810	0.032	¹⁴⁰ Nd	0.740	0.014
¹³³ Ce	0.810	0.022	¹⁴¹ Nd	0.750	0.014
¹³⁴ Ce	0.830	0.014	¹⁴² Nd	0.780	0.014
¹³⁵ Ce	0.840	0.014	¹⁴³ Nd	0.780	0.014
¹³⁶ Ce	0.840	0.014	¹⁴⁴ Nd	0.760	0.014
¹³⁷ Ce	0.860	0.014	¹⁴⁵ Nd	0.790	0.014
¹³⁸ Ce	0.860	0.014	¹⁴⁶ Nd	0.800	0.010
¹³⁹ Ce	0.850	0.014	¹⁴⁷ Nd	0.800	0.022
¹⁴⁰ Ce	0.860	0.014	¹⁴⁸ Nd	0.810	0.014
¹⁴¹ Ce	0.860	0.014	¹⁴⁹ Nd	0.820	0.014

TABLE VII: Velocities of spallation-fission fragments (in cm/ns) measured in the frame of the fissioning system for the reaction ²³⁸U(1A GeV)+d, with their total error.

nucleus	v_{fiss} (cm/ns)	ε_{tot} (cm/ns)	nucleus	v_{fiss} (cm/ns)	ε_{tot} (cm/ns)
¹⁴⁹ Nd	0.820	0.014	¹⁴⁸ Eu	0.690	0.041
¹⁵⁰ Nd	0.830	0.022	¹⁴⁹ Eu	0.700	0.040
¹⁵¹ Nd	0.830	0.022	¹⁵⁰ Eu	0.710	0.014
¹⁵² Nd	0.830	0.051	¹⁵¹ Eu	0.720	0.010
¹⁵³ Nd	0.810	0.041	¹⁵² Eu	0.710	0.014
¹⁵⁴ Nd	0.800	0.051	¹⁵³ Eu	0.720	0.014
¹⁵⁵ Nd	0.810	0.022	¹⁵⁴ Eu	0.750	0.040
¹⁵⁶ Nd	0.820	0.014	¹⁵⁵ Eu	0.750	0.022
¹³³ Pm	0.460	0.150	¹⁵⁶ Eu	0.750	0.010
¹³⁴ Pm	0.520	0.120	¹⁵⁷ Eu	0.750	0.041
¹³⁵ Pm	0.590	0.080	¹⁴⁹ Gd	0.640	0.051
¹³⁶ Pm	0.650	0.030	¹⁵⁰ Gd	0.640	0.041
¹³⁷ Pm	0.670	0.032	¹⁵¹ Gd	0.670	0.041
¹³⁸ Pm	0.700	0.028	¹⁵² Gd	0.690	0.010
¹³⁹ Pm	0.690	0.030	¹⁵³ Gd	0.670	0.022
¹⁴⁰ Pm	0.740	0.014	¹⁵⁴ Gd	0.700	0.041
¹⁴¹ Pm	0.760	0.014	¹⁵⁵ Gd	0.700	0.014
¹⁴² Pm	0.780	0.014	¹⁵⁶ Gd	0.710	0.014
¹⁴³ Pm	0.760	0.022	¹⁵⁷ Gd	0.720	0.014
¹⁴⁴ Pm	0.800	0.022	¹⁵⁸ Gd	0.710	0.032
¹⁴⁵ Pm	0.770	0.022	¹⁵⁹ Gd	0.710	0.022
¹⁴⁶ Pm	0.790	0.014	¹⁵⁴ Tb	0.680	0.032
¹⁴⁷ Pm	0.810	0.014	¹⁵⁵ Tb	0.690	0.030
¹⁴⁸ Pm	0.820	0.014	¹⁵⁶ Tb	0.700	0.010
¹⁴⁹ Pm	0.830	0.014	¹⁵⁷ Tb	0.700	0.010
¹⁵⁰ Pm	0.810	0.014	¹⁵⁸ Tb	0.690	0.050
¹⁵¹ Pm	0.820	0.014	¹⁵⁹ Tb	0.700	0.022
¹⁵² Pm	0.810	0.014	¹⁶⁰ Tb	0.700	0.020
¹⁵³ Pm	0.800	0.022	¹⁵⁶ Dy	0.660	0.061
¹⁵⁴ Pm	0.810	0.032	¹⁵⁷ Dy	0.660	0.051
¹⁵⁵ Pm	0.800	0.064	¹⁵⁸ Dy	0.650	0.032
¹⁵⁶ Pm	0.810	0.022	¹⁵⁹ Dy	0.670	0.010
¹⁵⁷ Pm	0.810	0.014	¹⁶⁰ Dy	0.660	0.051
¹⁵⁸ Pm	0.800	0.022	¹⁶¹ Dy	0.670	0.010
¹³⁹ Sm	0.600	0.050	¹⁶² Dy	0.670	0.060
¹⁴⁰ Sm	0.630	0.060			
¹⁴¹ Sm	0.650	0.032			
¹⁴² Sm	0.690	0.022			
¹⁴³ Sm	0.710	0.022			
¹⁴⁴ Sm	0.730	0.014			
¹⁴⁵ Sm	0.730	0.014			
¹⁴⁶ Sm	0.740	0.014			
¹⁴⁷ Sm	0.740	0.022			
¹⁴⁸ Sm	0.760	0.014			
¹⁴⁹ Sm	0.760	0.014			
¹⁵⁰ Sm	0.760	0.014			
¹⁵¹ Sm	0.760	0.014			
¹⁵² Sm	0.760	0.014			
¹⁵³ Sm	0.770	0.010			
¹⁵⁴ Sm	0.780	0.014			
¹⁵⁵ Sm	0.780	0.014			
¹⁵⁶ Sm	0.770	0.022			
¹⁵⁷ Sm	0.770	0.032			
¹⁵⁸ Sm	0.760	0.061			
¹⁵⁹ Sm	0.770	0.022			
¹⁶⁰ Sm	0.770	0.032			
¹⁴⁷ Eu	0.660	0.051			

TABLE VIII: Velocities of spallation-fission fragments (in cm/ns) measured in the frame of the fissioning system for the reaction ²³⁸U(1A GeV)+d, with their total error.

nucleus	σ (mb)	ε_{STAT} (mb)	ε_{SYST} (mb)	nucleus	σ (mb)	ε_{STAT} (mb)	ε_{SYST} (mb)
⁴⁹ V	0.159	0.003	0.021	⁶⁵ Ni	3.886	0.096	0.774
⁵⁰ V	0.770	0.013	0.099	⁶⁶ Ni	2.549	0.094	0.508
⁵¹ V	1.771	0.040	0.226	⁶⁷ Ni	1.745	0.087	0.348
⁵² V	1.618	0.059	0.091	⁶⁸ Ni	0.770	0.054	0.073
⁵³ V	1.470	0.074	0.083	⁶⁹ Ni	0.437	0.032	0.090
⁵⁴ V	0.973	0.078	0.123	⁷⁰ Ni	0.129	0.012	0.015
⁵⁵ V	0.411	0.053	0.054	⁵⁹ Cu	0.005	0.001	0.002
⁵⁶ V	0.219	0.037	0.032	⁶⁰ Cu	0.051	0.002	0.018
⁵⁷ V	0.019	0.003	0.002	⁶¹ Cu	0.505	0.007	0.125
⁵¹ Cr	0.375	0.005	0.059	⁶² Cu	1.714	0.015	0.379
⁵² Cr	1.333	0.017	0.202	⁶³ Cu	3.304	0.028	0.696
⁵³ Cr	2.373	0.042	0.361	⁶⁴ Cu	4.286	0.044	0.895
⁵⁴ Cr	2.485	0.061	0.377	⁶⁵ Cu	5.252	0.067	1.099
⁵⁵ Cr	2.044	0.082	0.310	⁶⁶ Cu	5.299	0.085	1.107
⁵⁶ Cr	1.331	0.074	0.202	⁶⁷ Cu	4.369	0.089	0.932
⁵⁷ Cr	0.551	0.053	0.037	⁶⁸ Cu	3.783	0.106	0.815
⁵⁸ Cr	0.281	0.037	0.020	⁶⁹ Cu	2.698	0.101	0.563
⁵⁹ Cr	0.068	0.010	0.007	⁷⁰ Cu	1.376	0.072	0.290
⁵³ Mn	0.536	0.007	0.094	⁷¹ Cu	1.094	0.062	0.104
⁵⁴ Mn	1.731	0.019	0.292	⁷² Cu	0.437	0.027	0.092
⁵⁵ Mn	2.683	0.037	0.452	⁷³ Cu	0.208	0.014	0.024
⁵⁶ Mn	2.725	0.056	0.463	⁷⁴ Cu	0.082	0.006	0.010
⁵⁷ Mn	2.833	0.082	0.478	⁶² Zn	0.020	0.001	0.004
⁵⁸ Mn	1.808	0.073	0.304	⁶³ Zn	0.308	0.004	0.069
⁵⁹ Mn	1.263	0.074	0.213	⁶⁴ Zn	1.356	0.012	0.293
⁶⁰ Mn	0.516	0.050	0.086	⁶⁵ Zn	3.039	0.024	0.659
⁶¹ Mn	0.283	0.029	0.048	⁶⁶ Zn	4.647	0.043	1.006
⁶² Mn	0.076	0.010	0.008	⁶⁷ Zn	6.196	0.070	1.339
⁵⁵ Fe	0.435	0.005	0.081	⁶⁸ Zn	6.349	0.088	1.373
⁵⁶ Fe	2.011	0.019	0.363	⁶⁹ Zn	5.969	0.102	1.291
⁵⁷ Fe	2.755	0.033	0.496	⁷⁰ Zn	5.093	0.111	1.145
⁵⁸ Fe	3.539	0.056	0.639	⁷¹ Zn	4.100	0.117	0.887
⁵⁹ Fe	2.779	0.064	0.530	⁷² Zn	2.448	0.095	0.532
⁶⁰ Fe	2.529	0.079	0.457	⁷³ Zn	1.227	0.063	0.120
⁶¹ Fe	1.864	0.084	0.336	⁷⁴ Zn	0.851	0.040	0.082
⁶² Fe	0.875	0.059	0.158	⁷⁵ Zn	0.421	0.022	0.092
⁶³ Fe	0.364	0.037	0.068	⁷⁶ Zn	0.119	0.006	0.027
⁶⁴ Fe	0.124	0.013	0.014	⁷⁷ Zn	0.051	0.003	0.012
⁶⁵ Fe	0.012	0.001	0.008	⁷⁸ Zn	0.018	0.001	0.005
⁵⁷ Co	0.112	0.001	0.023	⁶⁵ Ga	0.084	0.001	0.020
⁵⁸ Co	1.427	0.013	0.271	⁶⁶ Ga	0.723	0.007	0.177
⁵⁹ Co	3.057	0.030	0.588	⁶⁷ Ga	2.460	0.020	0.537
⁶⁰ Co	3.476	0.047	0.670	⁶⁸ Ga	4.325	0.039	0.923
⁶¹ Co	3.503	0.062	0.696	⁶⁹ Ga	6.319	0.066	1.349
⁶² Co	3.279	0.081	0.623	⁷⁰ Ga	7.708	0.094	1.654
⁶³ Co	2.533	0.083	0.500	⁷¹ Ga	8.182	0.115	1.749
⁶⁴ Co	1.800	0.087	0.342	⁷² Ga	6.917	0.121	1.501
⁶⁵ Co	0.846	0.062	0.163	⁷³ Ga	5.710	0.132	1.218
⁶⁶ Co	0.386	0.030	0.073	⁷⁴ Ga	3.891	0.120	0.831
⁶⁷ Co	0.266	0.023	0.052	⁷⁵ Ga	1.962	0.084	0.448
⁶⁸ Co	0.095	0.010	0.011	⁷⁶ Ga	1.663	0.065	0.159
⁶⁹ Co	0.042	0.005	0.006	⁷⁷ Ga	0.879	0.031	0.197
⁶⁰ Ni	0.567	0.005	0.114	⁷⁸ Ga	0.422	0.020	0.056
⁶¹ Ni	2.768	0.025	0.559	⁷⁹ Ga	0.291	0.011	0.037
⁶² Ni	3.370	0.037	0.673	⁸⁰ Ga	0.119	0.005	0.014
⁶³ Ni	4.156	0.063	0.830	⁶⁷ Ge	0.073	0.001	0.018
⁶⁴ Ni	3.916	0.075	0.816	⁶⁸ Ge	0.465	0.005	0.122

TABLE IX: Isotopic cross-sections of spallation-fission fragments of ²³⁸U(1A GeV)+d, with their statistical and systematic uncertainties.

nucleus	σ (mb)	ε_{STAT} (mb)	ε_{SYST} (mb)	nucleus	σ (mb)	ε_{STAT} (mb)	ε_{SYST} (mb)
⁶⁹ Ge	2.159	0.018	0.515	⁷⁸ Br	7.264	0.069	1.694
⁷⁰ Ge	3.997	0.033	0.903	⁷⁹ Br	10.398	0.100	2.416
⁷¹ Ge	6.243	0.061	1.379	⁸⁰ Br	12.284	0.124	2.852
⁷² Ge	8.447	0.090	1.868	⁸¹ Br	14.471	0.162	3.359
⁷³ Ge	9.475	0.118	2.091	⁸² Br	12.041	0.163	2.797
⁷⁴ Ge	8.801	0.128	1.969	⁸³ Br	10.773	0.174	2.484
⁷⁵ Ge	7.405	0.142	1.637	⁸⁴ Br	8.430	0.166	1.943
⁷⁶ Ge	5.796	0.134	1.305	⁸⁵ Br	5.936	0.153	0.705
⁷⁷ Ge	4.048	0.121	0.922	⁸⁶ Br	5.630	0.103	0.591
⁷⁸ Ge	2.438	0.088	0.254	⁸⁷ Br	5.336	0.078	0.548
⁷⁹ Ge	1.470	0.045	0.158	⁸⁸ Br	3.583	0.054	0.826
⁸⁰ Ge	1.255	0.034	0.277	⁸⁹ Br	2.128	0.032	0.492
⁸¹ Ge	0.712	0.020	0.158	⁹⁰ Br	0.791	0.014	0.181
⁸² Ge	0.450	0.012	0.100	⁹¹ Br	0.317	0.006	0.040
⁸³ Ge	0.151	0.004	0.020	⁹² Br	0.070	0.003	0.011
⁶⁹ As	0.053	0.001	0.013	⁷⁶ Kr	0.081	0.001	0.021
⁷⁰ As	0.481	0.006	0.109	⁷⁷ Kr	0.790	0.009	0.196
⁷¹ As	1.600	0.014	0.359	⁷⁸ Kr	2.056	0.018	0.482
⁷² As	3.586	0.030	0.801	⁷⁹ Kr	3.965	0.035	0.904
⁷³ As	6.194	0.056	1.381	⁸⁰ Kr	6.672	0.059	1.532
⁷⁴ As	7.688	0.079	1.715	⁸¹ Kr	10.931	0.099	2.492
⁷⁵ As	10.819	0.119	2.438	⁸² Kr	13.344	0.120	3.090
⁷⁶ As	10.065	0.131	2.246	⁸³ Kr	16.319	0.166	3.724
⁷⁷ As	10.433	0.157	2.328	⁸⁴ Kr	15.467	0.183	3.549
⁷⁸ As	7.783	0.151	1.738	⁸⁵ Kr	13.574	0.186	3.095
⁷⁹ As	5.915	0.138	1.374	⁸⁶ Kr	10.429	0.172	2.420
⁸⁰ As	4.191	0.138	0.430	⁸⁷ Kr	8.922	0.172	2.033
⁸¹ As	2.890	0.073	0.292	⁸⁸ Kr	8.752	0.173	0.917
⁸² As	2.235	0.050	0.222	⁸⁹ Kr	6.873	0.093	0.705
⁸³ As	1.688	0.038	0.377	⁹⁰ Kr	5.934	0.069	1.356
⁸⁴ As	0.434	0.010	0.097	⁹¹ Kr	3.999	0.055	0.912
⁸⁵ As	0.262	0.006	0.059	⁹² Kr	1.925	0.025	0.440
⁸⁶ As	0.131	0.004	0.017	⁹³ Kr	0.709	0.011	0.161
⁷¹ Se	0.031	0.001	0.009	⁹⁴ Kr	0.225	0.005	0.029
⁷² Se	0.329	0.004	0.085	⁹⁵ Kr	0.055	0.002	0.008
⁷³ Se	1.369	0.012	0.315	⁷⁸ Rb	0.037	0.001	0.009
⁷⁴ Se	3.053	0.025	0.703	⁷⁹ Rb	0.458	0.006	0.114
⁷⁵ Se	5.501	0.047	1.269	⁸⁰ Rb	1.572	0.015	0.376
⁷⁶ Se	7.932	0.075	1.828	⁸¹ Rb	3.455	0.029	0.810
⁷⁷ Se	10.228	0.108	2.358	⁸² Rb	6.654	0.058	1.521
⁷⁸ Se	12.497	0.136	2.879	⁸³ Rb	10.221	0.087	2.359
⁷⁹ Se	12.054	0.160	2.780	⁸⁴ Rb	14.326	0.123	3.316
⁸⁰ Se	10.764	0.167	2.481	⁸⁵ Rb	17.249	0.157	4.034
⁸¹ Se	8.513	0.162	1.963	⁸⁶ Rb	17.218	0.184	3.963
⁸² Se	6.492	0.149	1.498	⁸⁷ Rb	15.941	0.191	3.634
⁸³ Se	4.949	0.124	0.520	⁸⁸ Rb	12.739	0.185	2.926
⁸⁴ Se	4.022	0.073	0.429	⁸⁹ Rb	11.586	0.191	2.664
⁸⁵ Se	3.034	0.050	0.698	⁹⁰ Rb	9.864	0.211	1.040
⁸⁶ Se	2.200	0.040	0.509	⁹¹ Rb	8.898	0.129	0.944
⁸⁷ Se	1.016	0.018	0.234	⁹² Rb	7.201	0.088	0.753
⁸⁸ Se	0.463	0.009	0.060	⁹³ Rb	5.593	0.065	1.276
⁸⁹ Se	0.130	0.004	0.017	⁹⁴ Rb	3.166	0.044	0.722
⁷³ Br	0.012	0.000	0.003	⁹⁵ Rb	1.536	0.021	0.351
⁷⁴ Br	0.164	0.003	0.044	⁹⁶ Rb	0.510	0.009	0.116
⁷⁵ Br	0.962	0.010	0.236	⁹⁷ Rb	0.217	0.005	0.028
⁷⁶ Br	2.496	0.021	0.581	⁸⁰ Sr	0.030	0.001	0.008
⁷⁷ Br	4.120	0.035	0.949	⁸¹ Sr	0.324	0.005	0.106

TABLE X: Isotopic cross-sections of spallation-fission fragments of ²³⁸U(1A GeV)+d, with their statistical and systematic uncertainties.

nucleus	σ (mb)	ε_{STAT} (mb)	ε_{SYST} (mb)	nucleus	σ (mb)	ε_{STAT} (mb)	ε_{SYST} (mb)
^{82}Sr	0.864	0.009	0.225	^{101}Zr	5.516	0.061	0.579
^{83}Sr	2.878	0.026	0.673	^{102}Zr	3.289	0.040	0.757
^{84}Sr	5.304	0.045	1.258	^{103}Zr	1.150	0.017	0.266
^{85}Sr	9.982	0.085	2.306	^{104}Zr	0.414	0.008	0.095
^{86}Sr	13.913	0.114	3.182	^{105}Zr	0.093	0.003	0.013
^{87}Sr	17.897	0.149	4.076	^{87}Nb	0.106	0.002	0.033
^{88}Sr	18.938	0.188	4.369	^{88}Nb	0.445	0.006	0.125
^{89}Sr	18.306	0.200	4.197	^{89}Nb	1.422	0.015	0.332
^{90}Sr	15.109	0.195	3.445	^{90}Nb	3.462	0.031	0.811
^{91}Sr	13.863	0.203	3.162	^{91}Nb	7.335	0.060	1.688
^{92}Sr	12.004	0.200	2.739	^{92}Nb	10.170	0.084	2.395
^{93}Sr	10.979	0.202	1.175	^{93}Nb	13.121	0.109	3.029
^{94}Sr	10.720	0.125	1.122	^{94}Nb	15.875	0.137	3.678
^{95}Sr	7.233	0.076	1.658	^{95}Nb	19.255	0.182	4.458
^{96}Sr	5.207	0.060	1.187	^{96}Nb	18.550	0.188	4.284
^{97}Sr	2.715	0.033	0.619	^{97}Nb	17.958	0.199	4.131
^{98}Sr	1.344	0.018	0.306	^{98}Nb	16.911	0.211	3.923
^{99}Sr	0.343	0.007	0.044	^{99}Nb	16.133	0.217	3.718
^{100}Sr	0.116	0.003	0.015	^{100}Nb	14.169	0.236	1.502
^{82}Y	0.037	0.001	0.016	^{101}Nb	13.049	0.170	1.411
^{83}Y	0.200	0.003	0.072	^{102}Nb	8.380	0.092	0.973
^{84}Y	0.825	0.010	0.200	^{103}Nb	7.247	0.072	0.762
^{85}Y	2.260	0.021	0.585	^{104}Nb	3.629	0.046	0.382
^{86}Y	5.244	0.045	1.221	^{105}Nb	1.913	0.027	0.240
^{87}Y	8.730	0.073	2.020	^{106}Nb	0.634	0.012	0.079
^{88}Y	14.169	0.116	3.286	^{107}Nb	0.223	0.005	0.029
^{89}Y	17.153	0.138	4.025	^{89}Mo	0.065	0.002	0.018
^{90}Y	18.829	0.174	4.335	^{90}Mo	0.188	0.003	0.043
^{91}Y	19.189	0.199	4.433	^{91}Mo	1.132	0.012	0.314
^{92}Y	17.069	0.200	3.926	^{92}Mo	2.744	0.025	0.623
^{93}Y	15.452	0.204	3.576	^{93}Mo	5.860	0.052	1.351
^{94}Y	14.498	0.214	3.343	^{94}Mo	8.661	0.075	1.979
^{95}Y	11.265	0.211	1.178	^{95}Mo	12.761	0.106	2.951
^{96}Y	10.508	0.141	1.308	^{96}Mo	15.633	0.132	3.630
^{97}Y	9.141	0.101	1.016	^{97}Mo	18.477	0.169	4.249
^{98}Y	6.172	0.067	1.421	^{98}Mo	20.955	0.200	4.798
^{99}Y	4.739	0.056	1.090	^{99}Mo	21.074	0.213	4.799
^{100}Y	1.918	0.025	0.441	^{100}Mo	20.067	0.226	4.564
^{101}Y	0.763	0.011	0.176	^{101}Mo	18.476	0.225	4.224
^{102}Y	0.216	0.005	0.028	^{102}Mo	17.654	0.233	4.035
^{84}Zr	0.005	0.000	0.001	^{103}Mo	14.549	0.240	1.532
^{85}Zr	0.141	0.003	0.034	^{104}Mo	11.788	0.127	1.323
^{86}Zr	0.394	0.005	0.094	^{105}Mo	7.935	0.088	0.833
^{87}Zr	1.842	0.018	0.433	^{106}Mo	5.116	0.057	0.534
^{88}Zr	3.849	0.033	0.926	^{107}Mo	2.385	0.037	0.543
^{89}Zr	8.169	0.066	1.905	^{108}Mo	1.131	0.019	0.257
^{90}Zr	12.143	0.097	2.831	^{109}Mo	0.344	0.009	0.043
^{91}Zr	16.194	0.132	3.759	^{110}Mo	0.119	0.004	0.015
^{92}Zr	17.942	0.157	4.158	^{91}Tc	0.045	0.001	0.010
^{93}Zr	18.508	0.185	4.290	^{92}Tc	0.246	0.004	0.069
^{94}Zr	18.217	0.195	4.190	^{93}Tc	0.855	0.010	0.217
^{95}Zr	17.963	0.215	4.137	^{94}Tc	2.397	0.024	0.550
^{96}Zr	15.344	0.208	3.533	^{95}Tc	4.171	0.039	0.969
^{97}Zr	14.472	0.216	3.333	^{96}Tc	6.673	0.062	1.555
^{98}Zr	13.364	0.229	1.443	^{97}Tc	9.851	0.085	2.316
^{99}Zr	11.323	0.125	1.275	^{98}Tc	14.187	0.120	3.322
^{100}Zr	9.751	0.093	1.061	^{99}Tc	16.592	0.147	3.869

TABLE XI: Isotopic cross-sections of spallation-fission fragments of $^{238}U(1A \text{ GeV})+d$, with their statistical and systematic uncertainties.

nucleus	σ (mb)	ε_{STAT} (mb)	ε_{SYST} (mb)	nucleus	σ (mb)	ε_{STAT} (mb)	ε_{SYST} (mb)
^{100}Tc	19.663	0.185	4.508	^{118}Rh	0.070	0.003	0.010
^{101}Tc	20.179	0.197	4.609	^{98}Pd	0.002	0.000	0.000
^{102}Tc	18.649	0.200	4.313	^{99}Pd	0.191	0.004	0.054
^{103}Tc	21.048	0.231	4.807	^{100}Pd	0.727	0.010	0.175
^{104}Tc	17.382	0.224	3.960	^{101}Pd	1.862	0.022	0.485
^{105}Tc	16.068	0.224	3.658	^{102}Pd	3.455	0.037	0.854
^{106}Tc	11.778	0.170	1.345	^{103}Pd	5.816	0.057	1.394
^{107}Tc	10.211	0.111	1.119	^{104}Pd	8.667	0.080	2.125
^{108}Tc	5.921	0.067	0.618	^{105}Pd	11.983	0.108	2.835
^{109}Tc	3.708	0.045	0.387	^{106}Pd	14.443	0.134	3.394
^{110}Tc	1.718	0.029	0.211	^{107}Pd	17.980	0.167	4.203
^{111}Tc	0.751	0.014	0.093	^{108}Pd	19.669	0.194	4.581
^{93}Ru	0.020	0.001	0.005	^{109}Pd	19.397	0.198	4.516
^{94}Ru	0.026	0.001	0.008	^{110}Pd	18.034	0.198	4.195
^{95}Ru	0.583	0.008	0.136	^{111}Pd	17.787	0.212	4.109
^{96}Ru	1.588	0.018	0.394	^{112}Pd	17.657	0.215	4.071
^{97}Ru	3.631	0.037	0.872	^{113}Pd	14.161	0.202	2.715
^{98}Ru	5.619	0.053	1.327	^{114}Pd	10.619	0.139	1.200
^{99}Ru	8.850	0.080	2.063	^{115}Pd	7.617	0.092	0.918
^{100}Ru	12.199	0.104	2.846	^{116}Pd	4.432	0.053	0.509
^{101}Ru	16.912	0.148	3.880	^{117}Pd	2.208	0.032	0.233
^{102}Ru	18.222	0.165	4.239	^{118}Pd	1.174	0.024	0.124
^{103}Ru	19.590	0.189	4.497	^{119}Pd	0.460	0.011	0.057
^{104}Ru	19.782	0.198	4.625	^{120}Pd	0.136	0.005	0.019
^{105}Ru	19.489	0.211	4.523	^{100}Ag	0.012	0.000	0.002
^{106}Ru	19.459	0.229	4.438	^{101}Ag	0.076	0.002	0.019
^{107}Ru	17.368	0.224	3.954	^{102}Ag	0.471	0.007	0.113
^{108}Ru	15.723	0.232	1.703	^{103}Ag	1.229	0.016	0.282
^{109}Ru	10.755	0.134	1.212	^{104}Ag	2.326	0.026	0.574
^{110}Ru	8.438	0.096	0.963	^{105}Ag	3.769	0.040	0.969
^{111}Ru	4.769	0.057	0.497	^{106}Ag	6.369	0.062	1.491
^{112}Ru	2.682	0.039	0.334	^{107}Ag	9.757	0.092	2.382
^{113}Ru	1.017	0.021	0.125	^{108}Ag	12.643	0.120	2.950
^{114}Ru	0.437	0.011	0.053	^{109}Ag	15.100	0.143	3.501
^{115}Ru	0.122	0.004	0.015	^{110}Ag	16.290	0.165	3.756
^{96}Rh	0.030	0.001	0.013	^{111}Ag	18.021	0.183	4.161
^{97}Rh	0.268	0.004	0.101	^{112}Ag	17.368	0.192	3.998
^{98}Rh	1.005	0.012	0.240	^{113}Ag	17.722	0.203	4.065
^{99}Rh	2.145	0.023	0.540	^{114}Ag	16.645	0.205	3.794
^{100}Rh	4.265	0.043	0.981	^{115}Ag	15.805	0.194	3.593
^{101}Rh	6.882	0.064	1.586	^{116}Ag	12.694	0.194	1.481
^{102}Rh	10.234	0.092	2.376	^{117}Ag	9.756	0.115	1.180
^{103}Rh	14.602	0.126	3.401	^{118}Ag	6.062	0.074	0.685
^{104}Rh	17.064	0.157	3.942	^{119}Ag	3.861	0.048	0.410
^{105}Rh	20.457	0.189	4.716	^{120}Ag	1.616	0.026	0.174
^{106}Rh	19.515	0.196	4.499	^{121}Ag	0.809	0.017	0.098
^{107}Rh	20.153	0.206	4.611	^{122}Ag	0.264	0.008	0.034
^{108}Rh	18.780	0.215	4.298	^{103}Cd	0.019	0.001	0.005
^{109}Rh	16.913	0.202	4.612	^{104}Cd	0.254	0.005	0.061
^{110}Rh	15.025	0.200	1.576	^{105}Cd	0.843	0.012	0.280
^{111}Rh	13.508	0.197	1.593	^{106}Cd	1.709	0.021	0.462
^{112}Rh	10.379	0.120	1.238	^{107}Cd	3.123	0.034	0.781
^{113}Rh	6.707	0.076	0.795	^{108}Cd	4.959	0.052	1.283
^{114}Rh	3.412	0.044	0.359	^{109}Cd	7.643	0.076	1.795
^{115}Rh	1.745	0.030	0.215	^{110}Cd	9.638	0.095	2.276
^{116}Rh	0.685	0.015	0.083	^{111}Cd	12.226	0.120	2.817
^{117}Rh	0.251	0.007	0.032	^{112}Cd	14.366	0.145	3.283

TABLE XII: Isotopic cross-sections of spallation-fission fragments of $^{238}U(1A \text{ GeV})+d$, with their statistical and systematic uncertainties.

nucleus	σ (mb)	ε_{STAT} (mb)	ε_{SYST} (mb)	nucleus	σ (mb)	ε_{STAT} (mb)	ε_{SYST} (mb)
¹¹³ Cd	15.128	0.160	3.435	¹²⁶ Sn	3.669	0.049	0.413
¹¹⁴ Cd	15.777	0.171	3.600	¹²⁷ Sn	2.771	0.036	0.297
¹¹⁵ Cd	16.475	0.190	3.731	¹²⁸ Sn	1.820	0.026	0.196
¹¹⁶ Cd	16.740	0.203	3.761	¹²⁹ Sn	1.343	0.019	0.166
¹¹⁷ Cd	15.113	0.190	3.410	¹³⁰ Sn	1.169	0.014	0.144
¹¹⁸ Cd	10.934	0.156	2.513	¹³¹ Sn	0.733	0.008	0.091
¹¹⁹ Cd	10.290	0.153	1.115	¹³² Sn	0.442	0.006	0.055
¹²⁰ Cd	7.704	0.096	0.801	¹³³ Sn	0.012	0.000	0.004
¹²¹ Cd	4.621	0.059	0.562	¹⁰⁹ Sb	0.009	0.001	0.002
¹²² Cd	2.467	0.035	0.257	¹¹⁰ Sb	0.057	0.002	0.014
¹²³ Cd	1.327	0.026	0.140	¹¹¹ Sb	0.168	0.005	0.042
¹²⁴ Cd	0.510	0.012	0.062	¹¹² Sb	0.570	0.010	0.139
¹²⁵ Cd	0.203	0.006	0.025	¹¹³ Sb	1.178	0.017	0.295
¹²⁶ Cd	0.077	0.003	0.010	¹¹⁴ Sb	2.131	0.029	0.513
¹⁰⁵ In	0.013	0.000	0.003	¹¹⁵ Sb	3.333	0.040	0.842
¹⁰⁶ In	0.114	0.003	0.029	¹¹⁶ Sb	5.252	0.062	1.347
¹⁰⁷ In	0.452	0.008	0.118	¹¹⁷ Sb	6.393	0.077	1.650
¹⁰⁸ In	1.084	0.015	0.265	¹¹⁸ Sb	8.164	0.094	2.079
¹⁰⁹ In	2.088	0.025	0.503	¹¹⁹ Sb	9.653	0.114	2.350
¹¹⁰ In	3.877	0.042	0.975	¹²⁰ Sb	10.961	0.132	2.603
¹¹¹ In	5.395	0.055	1.385	¹²¹ Sb	11.627	0.143	2.747
¹¹² In	7.881	0.082	2.007	¹²² Sb	11.060	0.144	2.570
¹¹³ In	10.381	0.105	2.550	¹²³ Sb	11.611	0.156	2.688
¹¹⁴ In	12.574	0.130	2.968	¹²⁴ Sb	10.784	0.147	2.481
¹¹⁵ In	13.844	0.151	3.296	¹²⁵ Sb	9.661	0.134	2.221
¹¹⁶ In	15.444	0.171	3.592	¹²⁶ Sb	7.116	0.119	1.633
¹¹⁷ In	15.690	0.178	3.638	¹²⁷ Sb	6.640	0.104	0.716
¹¹⁸ In	14.787	0.183	3.407	¹²⁸ Sb	5.094	0.068	0.579
¹¹⁹ In	13.862	0.176	3.192	¹²⁹ Sb	3.929	0.047	0.423
¹²⁰ In	12.203	0.159	2.811	¹³⁰ Sb	3.203	0.037	0.344
¹²¹ In	10.288	0.166	1.109	¹³¹ Sb	3.059	0.034	0.703
¹²² In	8.373	0.117	0.947	¹³² Sb	2.227	0.023	0.274
¹²³ In	6.136	0.080	0.694	¹³³ Sb	1.720	0.017	0.212
¹²⁴ In	3.472	0.047	0.378	¹³⁴ Sb	0.096	0.001	0.026
¹²⁵ In	1.783	0.027	0.191	¹³⁵ Sb	0.029	0.001	0.009
¹²⁶ In	0.968	0.020	0.103	¹¹² Te	0.024	0.001	0.007
¹²⁷ In	0.548	0.012	0.068	¹¹³ Te	0.115	0.003	0.027
¹²⁸ In	0.038	0.001	0.009	¹¹⁴ Te	0.329	0.007	0.079
¹⁰⁷ Sn	0.008	0.001	0.002	¹¹⁵ Te	0.841	0.014	0.205
¹⁰⁸ Sn	0.034	0.001	0.009	¹¹⁶ Te	1.537	0.021	0.387
¹⁰⁹ Sn	0.281	0.006	0.064	¹¹⁷ Te	2.559	0.034	0.620
¹¹⁰ Sn	0.727	0.011	0.177	¹¹⁸ Te	3.488	0.046	0.910
¹¹¹ Sn	1.822	0.023	0.440	¹¹⁹ Te	4.839	0.063	1.246
¹¹² Sn	2.825	0.033	0.735	¹²⁰ Te	6.182	0.077	1.655
¹¹³ Sn	4.408	0.051	1.103	¹²¹ Te	6.582	0.086	1.664
¹¹⁴ Sn	6.598	0.073	1.689	¹²² Te	8.342	0.110	2.066
¹¹⁵ Sn	8.391	0.091	2.113	¹²³ Te	9.281	0.127	2.194
¹¹⁶ Sn	9.622	0.105	2.406	¹²⁴ Te	8.650	0.124	2.087
¹¹⁷ Sn	11.996	0.134	2.826	¹²⁵ Te	9.477	0.139	2.215
¹¹⁸ Sn	11.668	0.134	2.781	¹²⁶ Te	9.294	0.137	2.143
¹¹⁹ Sn	12.795	0.150	3.035	¹²⁷ Te	8.778	0.128	2.022
¹²⁰ Sn	12.038	0.156	2.769	¹²⁸ Te	6.930	0.106	0.761
¹²¹ Sn	12.193	0.159	2.814	¹²⁹ Te	6.561	0.122	0.717
¹²² Sn	11.054	0.146	2.555	¹³⁰ Te	5.171	0.071	0.557
¹²³ Sn	9.376	0.136	1.003	¹³¹ Te	4.740	0.057	0.549
¹²⁴ Sn	7.555	0.127	0.867	¹³² Te	4.721	0.047	0.531
¹²⁵ Sn	5.881	0.078	0.682	¹³³ Te	4.802	0.041	0.518

TABLE XIII: Isotopic cross-sections of spallation-fission fragments of ²³⁸U(1A GeV)+d, with their statistical and systematic uncertainties.

nucleus	σ (mb)	ε_{STAT} (mb)	ε_{SYST} (mb)	nucleus	σ (mb)	ε_{STAT} (mb)	ε_{SYST} (mb)
¹²⁰ Sb	10.961	0.132	2.603	¹²⁸ I	7.208	0.115	1.652
¹²¹ Sb	11.627	0.143	2.747	¹²⁹ I	7.704	0.117	1.782
¹²² Sb	11.060	0.144	2.570	¹³⁰ I	7.240	0.114	1.665
¹²³ Sb	11.611	0.156	2.688	¹³¹ I	5.971	0.106	0.644
¹²⁴ Sb	10.784	0.147	2.481	¹³² I	5.263	0.095	0.576
¹²⁵ Sb	9.661	0.134	2.221	¹³³ I	5.618	0.071	0.643
¹²⁶ Sb	7.116	0.119	1.633	¹³⁴ I	5.121	0.055	0.572
¹²⁷ Sb	6.640	0.104	0.716	¹³⁵ I	5.569	0.048	0.599
¹²⁸ Sb	5.094	0.068	0.579	¹³⁶ I	3.343	0.033	0.361
¹²⁹ Sb	3.929	0.047	0.423	¹³⁷ I	2.724	0.029	0.333
¹³⁰ Sb	3.203	0.037	0.344	¹³⁸ I	1.472	0.017	0.181
¹³¹ Sb	3.059	0.034	0.703	¹³⁹ I	0.139	0.002	0.034
¹³² Sb	2.227	0.023	0.274	¹⁴⁰ I	0.035	0.001	0.009
¹³³ Sb	1.720	0.017	0.212	¹⁴¹ I	0.011	0.000	0.003
¹³⁴ Sb	0.096	0.001	0.026	¹¹⁷ Xe	0.042	0.002	0.009
¹³⁵ Sb	0.029	0.001	0.009	¹¹⁸ Xe	0.131	0.004	0.030
¹¹² Te	0.024	0.001	0.007	¹¹⁹ Xe	0.403	0.009	0.101
¹¹³ Te	0.115	0.003	0.027	¹²⁰ Xe	0.920	0.015	0.222
¹¹⁴ Te	0.329	0.007	0.079	¹²¹ Xe	1.578	0.024	0.455
¹¹⁵ Te	0.841	0.014	0.205	¹²² Xe	2.034	0.030	0.581
¹¹⁶ Te	1.537	0.021	0.387	¹²³ Xe	2.824	0.043	0.797
¹¹⁷ Te	2.559	0.034	0.620	¹²⁴ Xe	3.999	0.059	1.067
¹¹⁸ Te	3.488	0.046	0.910	¹²⁵ Xe	5.219	0.076	1.345
¹¹⁹ Te	4.839	0.063	1.246	¹²⁶ Xe	4.898	0.074	1.363
¹²⁰ Te	6.182	0.077	1.655	¹²⁷ Xe	6.186	0.096	1.540
¹²¹ Te	6.582	0.086	1.664	¹²⁸ Xe	6.546	0.104	1.606
¹²² Te	8.342	0.110	2.066	¹²⁹ Xe	6.572	0.108	1.576
¹²³ Te	9.281	0.127	2.194	¹³⁰ Xe	6.786	0.114	1.612
¹²⁴ Te	8.650	0.124	2.087	¹³¹ Xe	6.535	0.109	1.541
¹²⁵ Te	9.477	0.139	2.215	¹³² Xe	6.230	0.105	1.432
¹²⁶ Te	9.294	0.137	2.143	¹³³ Xe	5.988	0.100	0.647
¹²⁷ Te	8.778	0.128	2.022	¹³⁴ Xe	5.450	0.107	0.601
¹²⁸ Te	6.930	0.106	0.761	¹³⁵ Xe	6.154	0.088	0.668
¹²⁹ Te	6.561	0.122	0.717	¹³⁶ Xe	6.596	0.073	0.755
¹³⁰ Te	5.171	0.071	0.557	¹³⁷ Xe	5.196	0.052	0.562
¹³¹ Te	4.740	0.057	0.549	¹³⁸ Xe	4.648	0.041	0.500
¹³² Te	4.721	0.047	0.531	¹³⁹ Xe	2.848	0.027	0.309
¹³³ Te	4.802	0.041	0.518	¹⁴⁰ Xe	2.108	0.022	0.256
¹³⁴ Te	4.534	0.040	0.557	¹⁴¹ Xe	1.030	0.013	0.128
¹³⁵ Te	2.009	0.020	0.247	¹⁴² Xe	0.071	0.001	0.019
¹³⁶ Te	0.202	0.002	0.051	¹⁴³ Xe	0.015	0.000	0.005
¹³⁷ Te	0.075	0.001	0.021	¹¹⁹ Cs	0.022	0.001	0.005
¹³⁸ Te	0.023	0.001	0.006	¹²⁰ Cs	0.088	0.003	0.021
¹³⁹ Te	0.004	0.000	0.001	¹²¹ Cs	0.258	0.006	0.065
¹¹⁵ I	0.064	0.003	0.013	¹²² Cs	0.630	0.012	0.153
¹¹⁶ I	0.183	0.005	0.047	¹²³ Cs	1.102	0.019	0.288
¹¹⁷ I	0.602	0.012	0.138	¹²⁴ Cs	1.756	0.028	0.467
¹¹⁸ I	1.204	0.020	0.312	¹²⁵ Cs	2.181	0.034	0.646
¹¹⁹ I	2.044	0.029	0.547	¹²⁶ Cs	3.203	0.050	0.937
¹²⁰ I	3.138	0.044	0.816	¹²⁷ Cs	3.385	0.054	1.040
¹²¹ I	4.010	0.056	1.004	¹²⁸ Cs	4.366	0.068	1.172
¹²² I	5.062	0.069	1.332	¹²⁹ Cs	4.306	0.070	1.169
¹²³ I	5.559	0.076	1.501	¹³⁰ Cs	5.149	0.088	1.244
¹²⁴ I	6.889	0.098	1.682	¹³¹ Cs	4.911	0.087	1.187
¹²⁵ I	7.440	0.109	1.822	¹³² Cs	4.952	0.090	1.166
¹²⁶ I	8.073	0.116	1.893	¹³³ Cs	4.893	0.089	1.131
¹²⁷ I	7.588	0.115	1.784	¹³⁴ Cs	4.654	0.084	1.065

TABLE XIV: Isotopic cross-sections of spallation-fission fragments of ²³⁸U(1A GeV)+d, with their statistical and systematic uncertainties.

nucleus	σ (mb)	ε_{STAT} (mb)	ε_{SYST} (mb)	nucleus	σ (mb)	ε_{STAT} (mb)	ε_{SYST} (mb)
¹³⁵ Cs	4.548	0.082	1.046	¹⁴¹ La	1.842	0.046	0.199
¹³⁶ Cs	4.116	0.077	0.455	¹⁴² La	2.120	0.058	0.232
¹³⁷ Cs	4.331	0.084	0.510	¹⁴³ La	2.106	0.042	0.230
¹³⁸ Cs	3.844	0.056	0.423	¹⁴⁴ La	1.937	0.034	0.217
¹³⁹ Cs	3.495	0.042	0.407	¹⁴⁵ La	2.074	0.028	0.248
¹⁴⁰ Cs	3.494	0.036	0.379	¹⁴⁶ La	1.200	0.017	0.144
¹⁴¹ Cs	3.116	0.029	0.338	¹⁴⁷ La	1.175	0.017	0.142
¹⁴² Cs	1.978	0.022	0.215	¹⁴⁸ La	0.458	0.010	0.056
¹⁴³ Cs	0.648	0.008	0.150	¹⁴⁹ La	0.059	0.002	0.014
¹⁴⁴ Cs	0.122	0.002	0.031	¹⁵⁰ La	0.013	0.001	0.004
¹⁴⁵ Cs	0.035	0.001	0.009	¹²⁶ Ce	0.023	0.001	0.006
¹⁴⁶ Cs	0.008	0.000	0.002	¹²⁷ Ce	0.086	0.003	0.020
¹²¹ Ba	0.016	0.001	0.004	¹²⁸ Ce	0.233	0.006	0.067
¹²² Ba	0.064	0.002	0.014	¹²⁹ Ce	0.602	0.013	0.153
¹²³ Ba	0.199	0.005	0.049	¹³⁰ Ce	0.794	0.016	0.307
¹²⁴ Ba	0.397	0.009	0.127	¹³¹ Ce	1.213	0.023	0.435
¹²⁵ Ba	0.822	0.016	0.225	¹³² Ce	1.181	0.024	0.638
¹²⁶ Ba	1.120	0.019	0.375	¹³³ Ce	1.627	0.032	0.646
¹²⁷ Ba	1.750	0.029	0.525	¹³⁴ Ce	1.716	0.036	0.707
¹²⁸ Ba	2.072	0.036	0.713	¹³⁵ Ce	1.887	0.040	0.718
¹²⁹ Ba	2.707	0.046	0.874	¹³⁶ Ce	1.793	0.041	0.675
¹³⁰ Ba	2.968	0.052	0.975	¹³⁷ Ce	2.280	0.054	0.602
¹³¹ Ba	3.550	0.063	0.975	¹³⁸ Ce	2.204	0.055	0.564
¹³² Ba	3.919	0.074	1.053	¹³⁹ Ce	2.127	0.054	0.508
¹³³ Ba	3.899	0.079	0.966	¹⁴⁰ Ce	2.109	0.054	0.499
¹³⁴ Ba	4.302	0.085	1.063	¹⁴¹ Ce	1.774	0.049	0.403
¹³⁵ Ba	3.729	0.077	0.873	¹⁴² Ce	1.470	0.046	0.336
¹³⁶ Ba	3.508	0.071	0.825	¹⁴³ Ce	1.369	0.039	0.358
¹³⁷ Ba	3.654	0.073	0.839	¹⁴⁴ Ce	1.251	0.038	0.137
¹³⁸ Ba	3.739	0.073	0.854	¹⁴⁵ Ce	1.506	0.046	0.169
¹³⁹ Ba	3.354	0.075	0.364	¹⁴⁶ Ce	1.496	0.033	0.167
¹⁴⁰ Ba	3.744	0.071	0.405	¹⁴⁷ Ce	1.373	0.027	0.151
¹⁴¹ Ba	3.508	0.049	0.381	¹⁴⁸ Ce	1.083	0.019	0.118
¹⁴² Ba	3.914	0.044	0.426	¹⁴⁹ Ce	0.763	0.015	0.083
¹⁴³ Ba	3.240	0.033	0.352	¹⁵⁰ Ce	0.463	0.011	0.056
¹⁴⁴ Ba	2.517	0.024	0.303	¹⁵¹ Ce	0.043	0.002	0.011
¹⁴⁵ Ba	0.755	0.010	0.174	¹⁵² Ce	0.015	0.001	0.005
¹⁴⁶ Ba	0.350	0.006	0.081	¹²⁸ Pr	0.023	0.002	0.005
¹⁴⁷ Ba	0.060	0.001	0.016	¹²⁹ Pr	0.052	0.002	0.013
¹⁴⁸ Ba	0.014	0.001	0.004	¹³⁰ Pr	0.129	0.004	0.052
¹²⁴ La	0.037	0.002	0.013	¹³¹ Pr	0.349	0.009	0.097
¹²⁵ La	0.110	0.003	0.027	¹³² Pr	0.623	0.014	0.203
¹²⁶ La	0.307	0.008	0.082	¹³³ Pr	0.882	0.019	0.353
¹²⁷ La	0.530	0.011	0.164	¹³⁴ Pr	1.145	0.024	0.454
¹²⁸ La	1.043	0.019	0.274	¹³⁵ Pr	0.962	0.022	0.582
¹²⁹ La	1.136	0.022	0.422	¹³⁶ Pr	1.330	0.030	0.571
¹³⁰ La	1.520	0.029	0.584	¹³⁷ Pr	1.160	0.029	0.584
¹³¹ La	1.771	0.035	0.611	¹³⁸ Pr	1.365	0.034	0.534
¹³² La	2.417	0.046	0.719	¹³⁹ Pr	1.419	0.038	0.474
¹³³ La	2.447	0.050	0.718	¹⁴⁰ Pr	1.859	0.049	0.484
¹³⁴ La	2.842	0.058	0.797	¹⁴¹ Pr	1.616	0.046	0.397
¹³⁵ La	2.759	0.061	0.731	¹⁴² Pr	1.447	0.044	0.342
¹³⁶ La	2.811	0.065	0.683	¹⁴³ Pr	1.290	0.042	0.305
¹³⁷ La	2.652	0.062	0.622	¹⁴⁴ Pr	1.209	0.042	0.278
¹³⁸ La	2.456	0.058	0.564	¹⁴⁵ Pr	1.192	0.042	0.273
¹³⁹ La	2.583	0.059	0.592	¹⁴⁶ Pr	1.046	0.036	0.114
¹⁴⁰ La	2.078	0.054	0.474	¹⁴⁷ Pr	0.835	0.033	0.092

TABLE XV: Isotopic cross-sections of spallation-fission fragments of ²³⁸U(1A GeV)+d, with their statistical and systematic uncertainties.

nucleus	σ (mb)	ε_{STAT} (mb)	ε_{SYST} (mb)	nucleus	σ (mb)	ε_{STAT} (mb)	ε_{SYST} (mb)
¹⁴⁸ Pr	0.794	0.029	0.088	¹⁵⁷ Pm	0.007	0.001	0.002
¹⁴⁹ Pr	0.819	0.023	0.091	¹⁵⁸ Pm	0.003	0.001	0.001
¹⁵⁰ Pr	0.725	0.020	0.080	¹⁴⁵ Sm	0.619	0.022	0.401
¹⁵¹ Pr	0.563	0.015	0.062	¹⁴⁶ Sm	0.448	0.017	0.356
¹⁵² Pr	0.210	0.007	0.023	¹⁴⁷ Sm	0.620	0.025	0.196
¹⁵³ Pr	0.099	0.005	0.012	¹⁴⁸ Sm	0.489	0.022	0.158
¹⁵⁴ Pr	0.008	0.001	0.003	¹⁴⁹ Sm	0.623	0.028	0.146
¹³¹ Nd	0.033	0.002	0.013	¹⁵⁰ Sm	0.420	0.020	0.109
¹³² Nd	0.080	0.003	0.038	¹⁵¹ Sm	0.336	0.018	0.084
¹³³ Nd	0.254	0.007	0.095	¹⁵² Sm	0.362	0.020	0.096
¹³⁴ Nd	0.450	0.011	0.210	¹⁵³ Sm	0.276	0.017	0.076
¹³⁵ Nd	0.662	0.016	0.346	¹⁵⁴ Sm	0.233	0.016	0.057
¹³⁶ Nd	0.906	0.021	0.458	¹⁵⁵ Sm	0.153	0.013	0.037
¹³⁷ Nd	0.988	0.023	0.593	¹⁵⁶ Sm	0.114	0.010	0.026
¹³⁸ Nd	0.992	0.025	0.656	¹⁵⁷ Sm	0.050	0.005	0.012
¹³⁹ Nd	1.061	0.028	0.560	¹⁵⁸ Sm	0.033	0.004	0.009
¹⁴⁰ Nd	0.867	0.026	0.598	¹⁵⁹ Sm	0.018	0.003	0.006
¹⁴¹ Nd	1.143	0.033	0.431	¹⁶⁰ Sm	0.009	0.002	0.004
¹⁴² Nd	1.145	0.035	0.363	¹⁴⁸ Eu	0.419	0.020	0.308
¹⁴³ Nd	1.182	0.039	0.297	¹⁴⁹ Eu	0.437	0.021	0.229
¹⁴⁴ Nd	1.034	0.037	0.244	¹⁵⁰ Eu	0.433	0.022	0.144
¹⁴⁵ Nd	0.951	0.035	0.220	¹⁵¹ Eu	0.336	0.019	0.107
¹⁴⁶ Nd	0.813	0.033	0.193	¹⁵² Eu	0.372	0.021	0.089
¹⁴⁷ Nd	0.696	0.031	0.168	¹⁵³ Eu	0.300	0.019	0.074
¹⁴⁸ Nd	0.666	0.030	0.163	¹⁵⁴ Eu	0.314	0.021	0.067
¹⁴⁹ Nd	0.568	0.025	0.130	¹⁵⁵ Eu	0.194	0.014	0.043
¹⁵⁰ Nd	0.481	0.023	0.052	¹⁵⁶ Eu	0.185	0.014	0.040
¹⁵¹ Nd	0.550	0.022	0.058	¹⁵⁷ Eu	0.087	0.009	0.019
¹⁵² Nd	0.358	0.014	0.038	¹⁵⁸ Eu	0.044	0.006	0.010
¹⁵³ Nd	0.262	0.011	0.029	¹⁵⁹ Eu	0.031	0.004	0.007
¹⁵⁴ Nd	0.141	0.007	0.020	¹⁶⁰ Eu	0.014	0.002	0.003
¹⁵⁵ Nd	0.076	0.005	0.010	¹⁵³ Gd	0.316	0.018	0.095
¹⁵⁶ Nd	0.005	0.001	0.001	¹⁵⁴ Gd	0.201	0.014	0.090
¹³³ Pm	0.027	0.002	0.007	¹⁵⁵ Gd	0.231	0.016	0.065
¹³⁴ Pm	0.069	0.003	0.018	¹⁵⁶ Gd	0.164	0.013	0.044
¹³⁵ Pm	0.181	0.006	0.060	¹⁵⁷ Gd	0.149	0.014	0.033
¹³⁶ Pm	0.410	0.012	0.142	¹⁵⁸ Gd	0.092	0.011	0.020
¹³⁷ Pm	0.516	0.014	0.262	¹⁵⁹ Gd	0.079	0.009	0.017
¹³⁸ Pm	0.909	0.022	0.393	¹⁶⁰ Gd	0.042	0.005	0.009
¹³⁹ Pm	0.634	0.018	0.555	¹⁶¹ Gd	0.017	0.003	0.004
¹⁴⁰ Pm	0.843	0.022	0.630	¹⁶² Gd	0.009	0.002	0.002
¹⁴¹ Pm	0.469	0.016	0.413	¹⁵² Tb	0.098	0.002	0.037
¹⁴² Pm	0.979	0.028	0.560	¹⁵³ Tb	0.131	0.004	0.038
¹⁴³ Pm	0.774	0.024	0.499	¹⁵⁴ Tb	0.168	0.005	0.037
¹⁴⁴ Pm	0.870	0.030	0.341	¹⁵⁵ Tb	0.165	0.006	0.023
¹⁴⁵ Pm	0.830	0.032	0.242	¹⁵⁶ Tb	0.183	0.011	0.020
¹⁴⁶ Pm	0.792	0.031	0.215	¹⁵⁷ Tb	0.184	0.008	0.017
¹⁴⁷ Pm	0.777	0.031	0.188	¹⁵⁸ Tb	0.162	0.009	0.014
¹⁴⁸ Pm	0.606	0.027	0.143	¹⁵⁹ Tb	0.100	0.009	0.009
¹⁴⁹ Pm	0.511	0.024	0.114	¹⁶⁰ Tb	0.067	0.005	0.006
¹⁵⁰ Pm	0.499	0.024	0.110	¹⁶¹ Tb	0.022	0.007	0.003
¹⁵¹ Pm	0.484	0.025	0.108	¹⁵⁴ Dy	0.041	0.001	0.019
¹⁵² Pm	0.336	0.020	0.074	¹⁵⁵ Dy	0.088	0.002	0.031
¹⁵³ Pm	0.326	0.020	0.037	¹⁵⁶ Dy	0.123	0.004	0.029
¹⁵⁴ Pm	0.254	0.016	0.028	¹⁵⁷ Dy	0.153	0.005	0.026
¹⁵⁵ Pm	0.172	0.012	0.018	¹⁵⁸ Dy	0.144	0.008	0.017
¹⁵⁶ Pm	0.094	0.007	0.010	¹⁵⁹ Dy	0.151	0.006	0.014

TABLE XVI: Isotopic cross-sections of spallation-fission fragments of ²³⁸U(1A GeV)+d, with their statistical and systematic uncertainties.

nucleus	σ (mb)	ε_{STAT} (mb)	ε_{SYST} (mb)
^{160}Dy	0.140	0.008	0.012
^{161}Dy	0.120	0.010	0.010
^{162}Dy	0.064	0.005	0.006
^{163}Dy	0.037	0.005	0.003
^{155}Ho	0.086	0.002	0.056
^{156}Ho	0.126	0.003	0.066
^{157}Ho	0.167	0.004	0.067
^{158}Ho	0.182	0.005	0.049
^{159}Ho	0.190	0.006	0.035
^{160}Ho	0.152	0.008	0.018
^{161}Ho	0.148	0.006	0.015
^{162}Ho	0.109	0.007	0.010
^{163}Ho	0.099	0.009	0.009
^{164}Ho	0.062	0.005	0.005
^{165}Ho	0.038	0.005	0.003
^{161}Er	0.043	0.001	0.012
^{162}Er	0.065	0.003	0.011
^{163}Er	0.082	0.003	0.010
^{164}Er	0.074	0.006	0.007
^{165}Er	0.059	0.009	0.005
^{166}Er	0.043	0.005	0.004
^{167}Er	0.023	0.004	0.002
^{165}Tm	0.005	0.000	0.001
^{166}Tm	0.032	0.002	0.004
^{167}Tm	0.049	0.005	0.004
^{168}Tm	0.047	0.006	0.004
^{169}Tm	0.028	0.004	0.003

TABLE XVII: Isotopic cross-sections of spallation-fission fragments of $^{238}U(1A\text{ GeV})+d$, with their statistical and systematic uncertainties.

- B. Mustapha, F. Rejmund, K.-H. Schmidt, C. Stéphan, L. Tassan-Got, and C. Volant, Nucl. Phys. A683 (2001) 513
- [14] F. Rejmund, B. Mustapha, P. Armbruster, J. Benlliure, M. Bernas, A. Boudard, J.P. Dufour, T. Enqvist, R. Legrain, S. Leray, M. Pravikoff, F. Rejmund, K.-H. Schmidt, C. Stéphan, J. Taïeb, L. Tassan-Got, C. Volant, and W. Wlazlo, Nucl. Phys. A 683 (2001) 540
- [15] T. Enqvist, W. Wlazlo, P. Armbruster, J. Benlliure, M. Bernas, A. Boudard, S. Czajkowski, R. Legrain, S. Leray, B. Mustapha, M. Pravikoff, F. Rejmund, K.-H. Schmidt, C. Stéphan, J. Taïeb, L. Tassan-Got, and C. Volant Nucl. Phys. A 686 (2001) 481
- [16] T. Enqvist, P. Armbruster, J. Benlliure, M. Bernas, A. Boudard, S. Czajkowski, R. Legrain, S. Leray, B. Mustapha, M. Pravikoff, F. Rejmund, K.-H. Schmidt, C. Stéphan, J. Taïeb, L. Tassan-Got, F. Vivès, C. Volant, and W. Wlazlo, Nucl. Phys. A 703 (2002) 435
- [17] J. Taïeb, K.-H. Schmidt, L. Tassan-Got, P. Armbruster, J. Benlliure, M. Bernas, A. Boudard, E. Casarejos, S. Czajkowski, T. Enqvist, R. Legrain, S. Leray, B. Mustapha, M. Pravikoff, F. Rejmund, C. Stéphan, C. Volant, and W. Wlazlo, Nucl. Phys. A724 (2003) 213
- [18] M. Bernas, P. Armbruster, J. Benlliure, A. Boudard, E. Casarejos, S. Czajkowski, T. Enqvist, R. Legrain, S. Leray, B. Mustapha, P. Napolitani, J. Pereira, F. Rejmund, M. V. Ricciardi, K.-H. Schmidt, C. Stéphan, J. Taïeb, L. Tassan-Got, C. Volant, Phys. Rev. Lett. 93 (2004) 212701
- [19] P. Armbruster, J. Benlliure, M. Bernas, A. Boudard, E. Casarejos, S. Czajkowski, T. Enqvist, S. Leray, P. Napolitani, J. Pereira, F. Rejmund, M.-V. Ricciardi, K.-H. Schmidt, C. Stéphan, J. Taïeb, L. Tassan-Got, and C. Volant, Phys. Rev. Lett. 93 (2004) 212701
- [20] C. Villagrasa, A. Boudard, J.E. Ducret, B. Fernández-Rodríguez, S. Leray, C. Volant, W. Wlazlo, P. Armbruster, T. Enqvist, F. Hammache, K. Helariutta, B. Jurado, M. V. Ricciardi, K.-H. Schmidt, K. Sümmerer, F. Vives, O. Yordanov, L. Audouin, L. Ferran, F. Rejmund, C. Stéphan, L. Tassan-Got, J. Benlliure, E. Casarejos, M. Fernández, J. Pereira, S. Czajkowski, D. Karamanis, M. Pravikoff, J. George, R.A. Mewaldt, N. Yanazak, A. Wiedenbeck, J. Connel, T. Faestermann, A. Heinz, and A. Junghans, AIP Conference Proceedings, Vol.769 (2005) 842
- [21] B. Fernández-Rodríguez, P. Armbruster, L. Audouin, J. Benlliure, M. Bernas, A. Boudard, E. Casarejos, S. Czajkowski, J.E. Ducret, T. Enqvist, B. Jurado, R. Legrain, S. Leray, B. Mustapha, J. Pereira, M. Pravikoff, F. Rejmund, M. V. Ricciardi, K.-H. Schmidt, C. Stéphan, J. Taïeb and L. Tassan-Got, and C. Volant, Nucl. Phys. A747 (2005) 227
- [22] M. Bernas, P. Armbruster, J. Benlliure, A. Boudard, E. Casarejos, S. Czajkowski, T. Enqvist, R. Legrain, S. Leray, B. Mustapha, P. Napolitani, J. Pereira, F. Rejmund, M. V. Ricciardi, K.-H. Schmidt, C. Stéphan, J. Taïeb, L. Tassan-Got, C. Volant, and W. Wlazlo, Nucl. Phys. A765 (2006) 197
- [23] M. V. Ricciardi, P. Armbruster, J. Benlliure, M. Bernas, A. Boudard, S. Czajkowski, T. Enqvist, A. Kelić, R. Legrain, S. Leray, B. Mustapha, J. Pereira, F. Rejmund, K.-H. Schmidt, C. Stéphan, L. Tassan-Got, C. Volant, and O. Yordanov, Phys. Rev. C73 (2006) 014607
- [24] C. Rubbia, J.A. Rubio, S. Buono, F. Carminati, N. Fiétier, J. Galvez, C. Gelés, Y. Kadi, R. Klapisch, P. Mandrillon, J.P. Revol, and Ch. Roche, Rep. CERN-AT/95-44(ET) (1995)
- [25] C.D.Bowman, Ann. Rev. Nucl. Part. Sci. 48 (1998) 505
- [26] <http://www.gsi.de>
- [27] M. Steiner, M. Blasche, H.-G. Clerc, H. Eickhoff, B. Franczak, H. Geissel, G. Münzenberg, K.-H. Schmidt, H. Stelzer, and K. Sümmerer, Nucl. Instr. and Meth. A312 (1992) 420
- [28] <http://www-aix.gsi.de/accelerator/sis.p14.html>
- [29] R. Anne, A. Lefol, G. Milleret, and R. Perret, Nucl. Instr. and Meth. 152 (1985) 395
- [30] B. Jurado, K.-H. Schmidt, and K.-H. Behr, Nucl. Instr. and Meth. A478 (2002) 493
- [31] Ph. Chesny, A. Forgeas, J.M. Gheller, G. Guiller, P. Pariset, L. Tassan-Got, P. Armbruster, K.-H. Behr, J. Benlliure, K. Burkard, A. Brünle, T. Enqvist, F. Farget, and K.-H. Schmidt, GSI-reports LNS/SSGD/93-73
- [32] H. Geissel, P. Armbruster, K.H. Behr, A. Brünle, K. Burkard, M. Chen, H. Folger, B. Franczak, H. Keller, O. Klepper, B. Langenbeck, F. Nickel, E. Pfeng, M. Pfützner, E. Roeckl, K. Rykaczewski, I. Schall, D. Scharadt, C. Scheidenberger, K.-H. Schmidt, A. Schröter, T. Schwab, K. Sümmerer, M. Weber, G. Münzenberg, T. Brohm, H.-G. Clerc, M. Fauer-

- bach, J.-J. Gaimard, A. Grewe, E. Hanelt, B. Knödler, M. Steiner, B. Voss, J. Weckenmann, C. Ziegler, A. Magel, H. Wollnik, J.P. Dufour, Y. Fujita, D.J. Vieira, and B. Sherrill, Nucl. Instr. and Meth. B70 (1992) 286
- [33] B. Voss, T. Brohm, H.-G. Clerc, A. Grewe, E. Hanelt, A. Heinz, M. de Jong, A. Junhans, W. Morawek, C. Röhl, S. Steinhäuser, C. Ziegler, K.-H. Schmidt, K.-H. Behr, H. Geissel, G. Münzenberg, F. Nickel, C. Scheidenberger, K. Sümmerer, A. Magel, and M. Pfützner Nucl. Instr. and Meth. A364 (1995) 150.
- [34] M. Pfützner, H. Geissel, G. Münzenberg, F. Nickel, C. Scheidenberger, K.-H. Schmidt, K. Sümmerer, T. Brohm, B. Voss, and H. Bichsel, Nucl. Instr. and Meth. B86 (1994) 213
- [35] H. Stelzer, Nucl. Instr. and Meth. A310 (1991) 103
- [36] J. Benlliure, J. Pereira, and K.-H. Schmidt, Nucl. Instr. Meth. A478 (2002) 493
- [37] J. Pereira, J. Benlliure, and K.-H. Schmidt, Nucl. Instr. and Meth. B204 (2003) 517
- [38] P. Karol, Phys. Rev. C11, 4 (1975) 1203
- [39] J.-J. Gaimard, and K.-H. Schmidt Nucl. Phys. A531 (1991) 709
- [40] P. Napolitani, L. Tassan-Got, P. Armbruster, and M. Bernas, Nucl. Phys. A727 (2003) 120
- [41] T. Enqvist, J. Benlliure, F. Farget, K.-H. Schmidt, P. Armbruster, M. Bernas, L. Tassan-Got, A. Boudard, R. Legrain, C. Volant, C. Böckstiegel, M. de Jong, and J.P. Dufour, Nucl. Phys. A658 (1999) 47
- [42] C. Scheidenberger, Th. Stöhlker, W.E. Meyerhof, H. Geissel, P.H. Mokler and B. Blank Nucl. Instr. Methods B142 (1998) 441
- [43] The GLOBAL program is available via anonymous ftp from borsu8.in2p3.fr in the directory */pub/nex/global/pc* and */pub/nex/global/vms_unix* for PCs and workstations, respectively
- [44] R. Serber, Phys. Rev. 72 (1947) 1114
- [45] J. Benlliure, A. Grewe, M. de Jong, K.-H. Schmidt, and S. Zhdanov Nucl. Phys. A 628 (1998) 458
- [46] B.D. Wilkins, E.P. Steinberg, and R.R. Chasman., Phys. Rev. C14 (1976) 1832
- [47] U. Brosa, S. Grossmann, and A. Müller, Phys. Rep. 197, 4 (1990) 167
- [48] The EXFOR entries are available at the OECD NEA Data Bank, <http://www.nea.fr/html/dbdata>
- [49] P.C. Stevenson, H.G. Hicks, W.E. Nervik, and D.R. Nethaway, Phys. Rev. 111 (1958) 886
- [50] A.V. Prokofiev, Nucl. Instr. and Meth. A463 (2001) 557
- [51] H. de Vries, W. de Jager, and C. de Vries, At. Data and Nucl. Data Tables 36 (1987) 495