

Seminar GANIL
Caen - June 2002

The Isospin Thermometer -
A New Tool to Determine
the Freeze-out Temperature
of Heavy Residues after
Relativistic Nucleon-Nucleon
Collisions

M. Valentina Ricciardi (thesis)

Alexandre Botvina


Timo Enqvist

Karl-Heinz Schmidt



GSI

INTRODUCTION

Signatures of the liquid-gas phase transition



investigation of
light clusters
(multifragmentation)



investigation of
heavy clusters
(fragmentation)

★ The formation of light clusters (multifragmentation) has widely been exploited to search for thermal instabilities of excited nuclei


★ Light clusters might be emitted by liquid and gaseous phase. - Heavy residues are clearly associated to the liquid phase.

★ The identification of heavy residues needs specific experimental tools.

OUTLOOK

- 1 Experiments at FRS of GSI
- 2 Results
- 3 Sequential decay or simultaneous break-up?
- 4 Idea behind the isospin thermometer
- 5 Comparison with a three stage model
- 6 Comparison with SMM calculations
- 7 Possible scenario of mid-peripheral high-energy nucleus-nucleus collisions
- 8 Conclusions

THE EXPERIMENT AT THE FRS AT GSI





1 A·GeV $^{238}\text{U} \rightarrow \text{Pb}$

Z from IC: $\Delta E \propto Z^2$

A/Z from time and position: $\frac{A}{Z} = \frac{e}{m_0} \frac{B\rho}{c\beta\gamma}$



T. Enqvist et al. / Nuclear Physics A 658 (1999) 47–66




velocity is calculated from $B\rho$:

$$\gamma v = B\rho \frac{Z \cdot e}{A \cdot m_0} \quad \text{very precise evaluation!}$$

DISCRIMINATION OF FISSION EVENTS



Systematic survey on residual nuclide production




6000 individual data points!

Basic data for

- EURISOL and GSI project
 - Intensities of secondary beams
- HINDAS
 - Nuclear data for incineration of nuclear waste


From electromagnetic-induced fission to fragmentation of ^{238}U



- Fission from excitations of GDR and nuclear collisions
- Fragmentation in high-energy nuclear collisions

Neutron excess reflects excitation energy induced.

EXPERIMENTAL RESULTS




Data: $^{238}\text{U} + ^{208}\text{Pb}$ (1 A GeV)
(Only fragmentation, fission discharged)

EPAX: a *systematics* of isotopic cross sections in projectile fragmentation

(K. Sümmerer, B. Blank, Phys. Rev. C (2000) 034607)

EPAX is based on the hypothesis of *limiting fragmentation*



Mean N/Z of fragments (fission discharged)



- stability line
- EPAX, projectile = Au
- EPAX, projectile = Fe
- 800 A·MeV Au + p - F.Rejmund NPA 683 (2001)
- 414 A·MeV Fe + p - W.R.Webber AJ 508 (1998)
- 1000 A·MeV U + Pb - T.Enqvist NPA 658 (1999)
- 1000 A·MeV U + Ti - M.V. Ricciardi's thesis (2002)

Why do some data agree with EPAX
and some deviate?


What can we learn from ALADIN data?



- Separation between multifragmentation and "spallation".
- $Z > 20$ is the heaviest fragment in the reaction.

Consolidated knowledge

- Lighter residues originate from more violent collisions
- More violent collisions \rightarrow larger excitation energy (ABRASION PICTURE)




SIMULTANEOUS
BREAK-UP

BOTH


SEQUENTIAL
DECAY

SIMULTANEOUS BREAK-UP




$N/Z \cong \text{constant}$

PHYSICAL REVIEW C 63 061601(R)





different
excitation
energies

SEQUENTIAL DECAY



IDEA BEHIND
LIMITING
FRAGMENTATION




For more violent collisions the evaporation starts at lower excitation energies !!!

PRINCIPLE OF THE ISOSPIN THERMOMETER

Simplifying hypotheses:

- only n-evaporation
- 15 MeV consumed for every evaporated n
- the evaporation stops when $\langle N_{\text{final}} \rangle / Z = 1.25$




$$\begin{aligned}
 E^* &= 15 \text{ MeV} \cdot \langle N_i - N_f \rangle \\
 \frac{\langle N_i - N_f \rangle}{Z} &\sim 0.34 \\
 E^* &= a T^2 \\
 a &\sim A/10 \text{ MeV}
 \end{aligned}
 \left. \vphantom{\begin{aligned} E^* &= 15 \text{ MeV} \cdot \langle N_i - N_f \rangle \\ \frac{\langle N_i - N_f \rangle}{Z} &\sim 0.34 \end{aligned}} \right\} \begin{aligned} E^*/Z &= \text{constant} \\ \text{or} \\ E^*/A &= \text{constant} \end{aligned}$$

$E^*/A = 0.1 T^2 \rightarrow T^2 = \text{constant}$


All pre-fragments start the evaporation cascade at a constant temperature!!!

ABRASION
+
SIMULTANEOUS BREAK-UP
+
SEQUENTIAL DECAY





COMPARISON WITH A THREE-STAGE MODEL

ABRASION / (BREAK-UP) / EVAPORATION
... complete but simplified...









COMPARISON WITH SMM CALCULATIONS

... not complete but more sophisticated...



- 1 A GeV ^{238}U on Ti measured at FRS
- 1 A GeV ^{238}U on Pb measured at FRS

A SHARP LIMITING TEMPERATURE?





- 1 A GeV ^{238}U on Ti measured at FRS
- 1 A GeV ^{238}U on Pb measured at FRS

Three-stage model

SMM (arbitray normalised)

POSSIBLE SCENARIO OF MID-PERIPHERAL HIGH-ENERGY NUCLEUS-NUCLEUS COLLISIONS



CONCLUSIONS

- ★ Some heavy residues produced in relativistic nucleus-nucleus collisions are unexpectedly neutron-rich
- ★ This neutron excess was interpreted as an indication for a simultaneous-break-up phase
- ★ The mean N/Z -ratio of the final elements can be used in combination with statistical-model codes in order to deduce the freeze-out temperature after break up ("isospin thermometer")
- ★ The average temperature of the break-up configuration at freeze out was determined to $T \approx 5 \text{ MeV}$
- ★ Consequence: The probability for an equilibrated compound nucleus to exist drops strongly above a limiting temperature of 5 MeV

<http://www-wnt.gsi.de/kschmidt/talks.htm>